



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Indican

Item No. 35417

CAS Registry No.: 487-60-5
Formal Name: 1H-indol-3-yl β-D-glucopyranoside
Synonyms: Indoxyl-β-D-glucoside, NSC 87517, Uroxanthin
MF: C₁₄H₁₇NO₆
FW: 295.3
Purity: ≥98%
UV/Vis.: λ_{max}: 224 nm
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years
Item Origin: Synthetic



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Indican is supplied as a solid. A stock solution may be made by dissolving the indican in the solvent of choice, which should be purged with an inert gas. Indican is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of indican in ethanol and DMSO is approximately 3 mg/ml and approximately 2 mg/ml in DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of indican can be prepared by directly dissolving the solid in aqueous buffers. The solubility of indican in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Indican is a glycoside and precursor in the synthesis of indigo that has been found in *I. tinctoria*.¹⁻³

References

1. Maugard, T., Enaud, E., de La Sayette, A., *et al.* β-glucosidase-catalyzed hydrolysis of indican from leaves of *Polygonum tinctorium*. *Biotechnol. Prog.* **18**(5), 1104-1108 (2002).
2. Kokubun, T., Edmonds, J.W., and John, P. Indoxyl derivatives in woad in relation to medieval indigo production. *Phytochemistry* **49**(1), 79-87 (1998).
3. Minami, Y., Kanafuji, T., and Miura, K. Purification and characterization of a β-glucosidase from *Polygonum tinctorium*, which catalyzes preferentially the hydrolysis of indican. *Biosci. Biotech. Biochem.* **60**(1), 147-149 (1996).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/23/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM