



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## IPG-2 TMA

Item No. 35533

**CAS Registry No.:** 1369302-21-5  
**Formal Name:** 4,5-dichloro-6-hydroxy-9-[3-methoxy-4-[16-(2-methoxy-4-methylphenyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadec-7-yl]phenyl]-3-oxo-3H-xanthene-2,7-dipropanoate, N,N,N-trimethyl-methanaminium (1:3)

**Synonyms:** APG-2, Asante Potassium Green-2, ION Potassium Green-2, IPG-2

**MF:**  $C_{46}H_{49}Cl_2N_2O_{13} \cdot 3C_4H_{12}N$

**FW:** 1,131.2

**Purity:**  $\geq 90\%$

**Ex./Em. Max:** 525/545 nm


**Supplied as:** A solid

**Storage:**  $-20^\circ C$

**Stability:**  $\geq 4$  years

**Special Conditions:** Protect from light and moisture.

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



### Laboratory Procedures

IPG-2 TMA is supplied as a solid. A stock solution may be made by dissolving the IPG-2 TMA in the solvent of choice, which should be purged with an inert gas. IPG-2 TMA is soluble in organic solvents such as methanol. It is also soluble in water. We do not recommend storing the aqueous solution for more than one day.

### Description

IPG-2 TMA is a cell-impermeable fluorescent potassium indicator. It binds to potassium ( $K_d = 18$  mM) and displays excitation/emission maxima of 525/545 nm, respectively. IPG-2 TMA can be used in lipid membrane-free systems or liposomes or can be introduced into cells by electroporation, microinjection, or other methods. IPG-2 is also available in a cell-permeable form (Item No. 35540).

WARNING  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/13/2022

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM