



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



ISG15 (mature form) (human, recombinant)

Item No. 35599

Overview and Properties

Synonyms: Interferon-induced 15 kDa Protein, Interferon-induced 17 kDa Protein, Interferon-stimulated Gene 15, IP17, Ubiquitin Cross-reactive Protein, Ubiquitin-like Protein ISG15, UCRP

Source: Recombinant human ISG15 expressed in *E. coli*

Amino Acids: 1-157

Uniprot No.: P05161

Molecular Weight: 17.2 kDa

Storage: -80°C (as supplied)


Stability: ≥1 year

Purity: ≥95% estimated by SDS-PAGE

Supplied in: Lyophilized from sterile 50 mM Tris, pH 8.0

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Image



Lane 1: MW Markers
Lane 2: ISG15 (mature form)

SDS-PAGE Analysis of ISG15 (mature form). This protein has a calculated molecular weight of 17.2 kDa. It has an apparent molecular weight of approximately 15 kDa by SDS-PAGE under reducing conditions.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/16/2021

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Interferon-stimulated gene 15 (ISG15) is a ubiquitin-like (Ubl) protein found only in vertebrates.^{1,2} The mature form of ISG15 is composed of an N-terminal Ubl and a C-terminal Ubl with an LRLRGG conjugation sequence at the C-terminus.² It is expressed in lymphoid cells, epithelial cells, neurons, and skeletal and smooth muscle, and its expression is induced by type I interferon signaling in response to viral or bacterial infection but is also induced in response to the aging process and certain diseases.¹⁻³ ISG15 is primarily localized intracellularly conjugated to lysine residues in newly synthesized proteins via ISGylation, a process that can enhance antiviral responses.¹ It also exists in a free form that can be secreted by a variety of cells, including fibroblasts, monocytes, and lymphocytes, and binds to the adhesion molecule lymphocyte function-associated antigen-1 (LFA-1) on leukocytes to increase the secretion of IFN- γ . ISG15 can ISGylate viral proteins to prevent viral replication and inhibit the function of viral proteins, however ISGylation on host proteins can be reversed by certain viruses, including severe acute respiratory coronavirus 2 (SARS-CoV-2), leading to suppression of the host innate immune response.^{1,4,5} Patients deficient in functional ISG15 due to inherited mutations in *ISG15* are more resistant to viral infection, but have an increased susceptibility to mycobacterial infection, while *Isg15* knockout mice are more susceptible to viral infection, indicating species-specific differences.^{6,7} Cayman's ISG15 (mature form) (human, recombinant) protein consists of 157 amino acids and has a calculated molecular weight of 17.2 kDa. By SDS-PAGE, under reducing conditions, the apparent molecular mass of the protein is 15 kDa.

References

1. Dzimianski, J.V., Scholte, F.E.M., Bergeron, É., *et al.* ISG15: It's complicated. *J. Mol. Biol.* **431(21)**, 4203-4216 (2019).
2. Zhang, D. and Zhang, D.-E. Interferon-stimulated gene 15 and the protein ISGylation system. *J. Interferon Cytokine Res.* **31(1)**, 119-130 (2011).
3. Lowe, J., McDermott, H., Loeb, K., *et al.* Immunohistochemical localization of ubiquitin cross-reactive protein in human tissues. *J. Pathol.* **177(2)**, 163-169 (1995).
4. Freitas, B.T., Durie, I.A., Murray, J., *et al.* Characterization and noncovalent inhibition of the deubiquitinase and deISGylase activity of SARS-CoV-2 papain-like protease. *ACS Infect. Dis.* **6(8)**, 2099-2109 (2020).
5. Shin, D., Mukherjee, R., Grewe, D., *et al.* Papain-like protease regulates SARS-CoV-2 viral spread and innate immunity. *Nature* **587(7835)**, 657-662 (2020).
6. Speer, S.D., Li, Z., Buta, S., *et al.* ISG15 deficiency and increased viral resistance in humans but not mice. *Nat. Commun.* **7**, 11496 (2016).
7. Bogunovic, D., Byun, M., Durfee, L.A., *et al.* Mycobacterial disease and impaired IFN- γ immunity in humans with inherited ISG15 deficiency. *Science* **337(6102)**, 1684-1688 (2012).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM