



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Mitochonic Acid 5

Item No. 35702

CAS Registry No.: 1354707-41-7
Formal Name: α -[2-(2,4-difluorophenyl)-2-oxoethyl]-1H-indole-3-acetic acid
Synonym: MA-5
MF: C₁₈H₁₃F₂NO₃
FW: 329.3
Purity: \geq 98%
UV/Vis.: λ_{max} : 220 nm
Supplied as: A solid
Storage: -20°C
Stability: \geq 4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Mitochonic acid 5 is supplied as a solid. A stock solution may be made by dissolving the mitochonic acid 5 in the solvent of choice, which should be purged with an inert gas. Mitochonic acid 5 is soluble in the organic solvent DMSO at a concentration of approximately 1 mg/ml. Mitochonic acid 5 is also slightly soluble in dimethyl formamide.

Description

Mitochonic acid 5 is an inhibitor of mitochondrial dysfunction.¹ It increases intracellular ATP levels in Hep3B cells when used at a concentration of 3 μ M. Mitochonic acid 5 (5 μ M) rescues LPS-induced apoptosis and decreases in the mitochondrial membrane potential in BV-2 microglial cells.² It protects against decreases in cell viability induced by L-buthionine-(S,R)-sulfoximine (BSO; Item No. 14484) in fibroblasts isolated from patients with various mitochondrial diseases, including Leigh syndrome and Leber's hereditary optic neuropathy (EC₅₀s = 2.3-5.1 μ M), in an electron transport chain-independent manner.¹ Mitochonic acid 5 (50 mg/kg) reduces plasma levels of creatinine and decreases acute renal tubular necrosis in a mouse model of bilateral retroperitoneal renal ischemia-reperfusion injury.³

References

1. Suzuki, T., Yamaguchi, H., Kikusato, M., *et al.* Mitochonic acid 5 (MA-5), a derivative of the plant hormone indole-3-acetic acid, improves survival of fibroblasts from patients with mitochondrial diseases. *Tohoku J. Exp. Med.* **236**(3), 225-232 (2015).
2. Tan, J., Chen, S.-X., Lei, Q.-Y., *et al.* Mitochonic acid 5 regulates mitofusin 2 to protect microglia. *Neural Regen. Res.* **16**(9), 1813-1820 (2021).
3. Suzuki, T., Yamaguchi, H., Kikusato, M., *et al.* Mitochonic acid 5 binds mitochondria and ameliorates renal tubular and cardiac myocyte damage. *J. Am. Soc. Nephrol.* **27**(7), 1925-1932 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/04/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM