



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



3-chloro-L-Tyrosine

Item No. 35753

CAS Registry No.: 7423-93-0
Synonyms: 3-Chlorotyrosine, 3-CT
MF: C₉H₁₀ClNO₃
FW: 215.6
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

3-chloro-L-Tyrosine is supplied as a solid. A stock solution may be made by dissolving the 3-chloro-L-tyrosine in the solvent of choice, which should be purged with an inert gas. 3-chloro-L-Tyrosine is slightly soluble in ethanol and DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 3-chloro-L-tyrosine can be prepared by directly dissolving the solid in aqueous buffers. The solubility of 3-chloro-L-tyrosine in PBS (pH 7.2) is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

3-chloro-L-Tyrosine is a derivative of L-tyrosine.¹ It is formed from a reaction between the myeloperoxidase (MPO) product hypochlorous acid and L-tyrosine and has been used as a biomarker of oxidative damage induced by MPO. Plasma and atherosclerotic plaque levels of 3-chloro-L-tyrosine are increased in patients with various cardiovascular diseases.² Plasma levels of 3-chloro-L-tyrosine are also increased in patients with colorectal cancer. 3-chloro-L-Tyrosine is also formed in blood upon chlorine gas exposure and has been used as an indicator of chlorine poisoning during autopsy.³

References

1. Feeney, M.B. and Schöneich, C. Tyrosine modifications in aging. *Antioxid. Redox Signal.* **17(11)**, 1571-1579 (2012).
2. Fleszar, M.G., Fortuna, P., Zawadzki, M., *et al.* Simultaneous LC-MS/MS-based quantification of free 3-nitro-L-tyrosine, 3-chloro-L-tyrosine, and 3-bromo-L-tyrosine in plasma of colorectal cancer patients during early postoperative period. *Molecules* **25(21)**, 5158 (2020).
3. Nishio, T., Toukairin, Y., Hoshi, T., *et al.* Determination of 3-chloro-L-tyrosine as a novel indicator of chlorine poisoning utilizing gas chromatography-mass spectrometric analysis. *Leg. Med. (Tokyo)* **47**, 101782 (2020).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/13/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM