



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Fluo-2 AM

Item No. 35758

CAS Registry No.: 1070771-36-6
Formal Name: N-[2-[(acetyloxy)methoxy]-2-oxoethyl]-N-[4-[6-[(acetyloxy)methoxy]-3-oxo-3H-xanthen-9-yl]-2-[2-[2-[bis[2-[(acetyloxy)methoxy]-2-oxoethyl]amino]-5-methylphenoxy]ethoxy]phenyl]-glycine (acetyloxy)methyl ester

Synonyms: Fluo-2 Acetoxymethyl ester, Fluo-2 HA, Fluo-2 High Affinity

MF: C₅₁H₅₂N₂O₂₃

FW: 1,061.0

Purity: ≥95%

UV/Vis.: λ_{max}: 254 nm

Ex./Em. Max: 490/515 nm


Supplied as: A solid

Storage: -20°C

Stability: ≥4 years

Special Conditions: Protect from light and moisture

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Description

Fluo-2 AM is a cell-permeable fluorescent calcium indicator.^{1,2} It has been used to measure the activity of the neuropeptide Y₄ receptor, as well as to detect intracellular calcium in tetrandine-stimulated primary rabbit corpus cavernosum smooth muscle cells. It binds to calcium (K_d = 290 nM) and displays excitation/emission maxima of 490/515 nm, respectively. Fluo-2 AM is also available in a cell-impermeable form (Item No. 35764).

Assay Protocol

Note: Allow all reagents to warm to room temperature before proceeding.

1. Add 10 ml of assay buffer to a 15 or 50 ml conical tube.

Note 1: HEPES-buffered Hank's balanced salt solution (HBSS), pH 7.2-7.4, is recommended, although other buffers can be used.

2. Add 100 μl of a 100X Pluronic™ F-127 solution (1-50% w/v) to the conical tube*. Pluronic™ F-127 is a biocompatible surfactant used to ensure equitable dye distribution and cellular loading.
 - a. Optional: Add 100 μl of 2 mM probenecid stock solution to the conical tube. Probenecid (Item No. 14981) is an anion transport inhibitor used to improve intracellular dye retention. Use of probenecid is recommended, but not required, for all cell types and dyes.

*Final working concentration of Pluronic™ F-127 should be between 0.01 and 0.5% w/v. User should optimize the concentration of Pluronic™ F-127 to suit experimental requirements.

Note 2: Probenecid is an inhibitor or agonist of multiple ion channels and may have undesirable cellular effects that could affect dye performance.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/13/2022

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



3. Vortex conical tube briefly to mix.
4. Dissolve Fluo-2 AM in 25 μ l of DMSO and vortex dye tube briefly to mix.
5. Centrifuge dye tube briefly to collect all contents at the tube bottom.
6. Add entire contents of dye tube to the conical tube containing the assay buffer solution to make the dye loading solution.
7. Vortex conical tube briefly to mix.

Note 3: The dye loading solution should be used within two hours for best results.

8. Remove cell culture medium and add dye loading solution. Recommended volumes are:
 - a. 35 mm dish or 6-well plate: 1.5 ml/dish or well
 - b. 96-well plate: 100 μ l/well
 - c. 384-well plate: 20 μ l/well

Note 4: To prevent cell detachment or if using suspension cells, the dye loading solution can be added directly to the media-containing wells. User must double the component concentrations to achieve the same final concentration of all reagents.

9. Incubate cells with the dye loading solution at 37°C for 60 minutes.
10. Read fluorescence using a plate reader at excitation and emission wavelengths of 490 and 515 nm, respectively.
Or
Image using a fluorescence microscope with filters for GFP or fluorescein.

References

1. Sliwoski, G., Schubert, M., Stichel, J., *et al.* Discovery of small-molecule modulators of the human Y_4 receptor. *PLoS One* **11(6)**, e0157146 (2016).
2. Liu, J.-H., Chen, J., Wang, T., *et al.* Effects of tetrandrine on cytosolic free calcium concentration in corpus cavernosum smooth muscle cells of rabbits. *Asian J. Androl.* **8(4)**, 405-409 (2006).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM