



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



rac-Hesperetin-d₃ Item No. 35802

CAS Registry No.: 1346605-26-2
Formal Name: 5,7-dihydroxy-2-(3-hydroxy-4-(methoxy-d₃)phenyl)chroman-4-one
Synonym: (±)-3',5,7-Trihydroxy-4'-methoxyflavanone-d₃
MF: C₁₆H₁₁D₃O₆
FW: 305.3
Chemical Purity: ≥95% (Hesperetin)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₃); ≤1% d₀
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years
Item Origin: Synthetic



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

rac-Hesperetin-d₃ is intended for use as an internal standard for the quantification of hesperetin (Item No. 10006084) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

rac-Hesperetin-d₃ is supplied as a solid. A stock solution may be made by dissolving the *rac*-hesperetin-d₃ in the solvent of choice, which should be purged with an inert gas. *rac*-Hesperetin-d₃ is slightly soluble in methanol.

Description

rac-Hesperetin is a racemic mixture of the flavanones (S)-hesperetin (Item No. 10006084) and (R)-hesperetin. *rac*-Hesperetin (1 mM) inhibits heat-induced advanced glycation end product formation between BSA and glucose by 56.7%.¹ It also inhibits α-glucosidase and protein tyrosine phosphatase 1B (PTP1B; IC₅₀s = 15.28 and 20.16 μM, respectively).² *rac*-Hesperetin (100 μM) protects against hydrogen peroxide-induced cytotoxicity in MIN6 pancreatic beta cells by 72.27%.

References

1. Li, D., Mitsuhashi, S., and Ubukata, M. Protective effects of hesperidin derivatives and their stereoisomers against advanced glycation end-products formation. *Pharm. Biol.* **50**(12), 1531-1535 (2012).
2. Xu, J., Wang, X., Yue, J., et al. Polyphenols from acorn leaves (*Quercus liaotungensis*) protect pancreatic beta cells and their inhibitory activity against α-glucosidase and protein tyrosine phosphatase 1B. *Molecules* **23**(9), 2167 (2018).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/13/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM