



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



O-desethyl Dapagliflozin

Item No. 36022

CAS Registry No.: 864070-37-1
Formal Name: (1S)-1,5-anhydro-1-C-[4-chloro-3-[(4-hydroxyphenyl)methyl]phenyl]-D-glucitol
MF: C₁₉H₂₁ClO₆
FW: 380.8
Purity: ≥98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

O-desethyl Dapagliflozin is supplied as a crystalline solid. A stock solution may be made by dissolving the O-desethyl dapagliflozin in the solvent of choice, which should be purged with an inert gas. O-desethyl Dapagliflozin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of O-desethyl dapagliflozin in these solvents is approximately 30 mg/ml.

O-desethyl Dapagliflozin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, O-desethyl dapagliflozin should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. O-desethyl Dapagliflozin has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

O-desethyl Dapagliflozin is an inhibitor of sodium-glucose cotransporter 2 (SGLT2; IC₅₀ = 9.4 nM) and active metabolite of dapagliflozin (Item No. 11574).^{1,2} It is selective for SGLT2 over SGLT1 (IC₅₀ = 970 nM).¹

References

- Xu, B., Feng, Y., Cheng, H., *et al.* C-aryl glucosides substituted at the 4'-position as potent and selective renal sodium-dependent glucose co-transporter 2 (SGLT2) inhibitors for the treatment of type 2 diabetes. *Bioorg. Med. Chem. Lett.* **21(15)**, 4465-4470 (2011).
- Xu, G., Lv, B., Roberge, J.Y., *et al.* Design, synthesis, and biological evaluation of deuterated C-aryl glycoside as a potent and long-acting renal sodium-dependent glucose cotransporter 2 inhibitor for the treatment of type 2 diabetes. *J. Med. Chem.* **57(4)**, 1236-1251 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/11/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM