



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Dansyl Amide

Item No. 36149

CAS Registry No.: 1431-39-6
Formal Name: 5-(dimethylamino)-1-naphthalenesulfonamide
Synonym: DNSA
MF: C₁₂H₁₄N₂O₂S
FW: 250.3
Purity: ≥98%
Ex./Em. Max: 280/470 nm
UV/Vis.: λ_{max}: 216, 249 nm
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Dansyl amide is supplied as a solid. A stock solution may be made by dissolving the dansyl amide in the solvent of choice, which should be purged with an inert gas. Dansyl amide is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of dansyl amide in these solvents is approximately 1 and 2 mg/ml, respectively.

Description

Dansyl amide is a fluorescent probe for albumin and CAII.^{1,2} It binds to serum albumin and carbonic anhydrase II (CAII; K_ds = 7.57 and 0.2 μM, respectively) and has been used to determine ligand affinities for these proteins. Dansyl amide binding to CAII results in an increase in fluorescence due to FRET from nearby active site tryptophan residues and displays excitation/emission maxima of 280/470 nm, respectively, when bound to CAII and a decrease in fluorescence can be used to quantify CAII ligand binding.²

References

1. Epps, D.E., Raub, T.J., and Kézdy, F.J. A general, wide-range spectrofluorometric method for measuring the site-specific affinities of drugs toward human serum albumin. *Anal. Biochem.* **227**(2), 342-350 (1995).
2. Wang, S.C. and Zamble, D.B. Fluorescence analysis of sulfonamide binding to carbonic anhydrase. *Biochem. Mol. Biol. Educ.* **34**(5), 364-368 (2006).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM