

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Koumine

Item No. 36210

CAS Registry No.: 1358-76-5

Formal Name: (3S,4S,4aR,6R,11bS,11cS)-11c-ethenyl-

1,2,3,4,4a,5,6,11c-octahydro-2-methyl-6,4-(epoxymethano)-3,11b-methano-

11bH-pyrido[4,3-c]carbazole


MF: $C_{20}H_{22}N_2O$ FW: 306.4 **Purity:** ≥98%

 λ_{max} : 215, 221 nm UV/Vis.:

Supplied as: A solid -20°C Storage: Stability: ≥4 years

Item No: Plant/Gelsemium elegans

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Koumine is supplied as a solid. A stock solution may be made by dissolving the koumine in the solvent of choice, which should be purged with an inert gas. Koumine is soluble in DMSO.

Description

Koumine is an alkaloid that has been found in Gelsemium elegans and has diverse biological activities. 1-4 It inhibits the proliferation of MCF-7 breast cancer cells (IC $_{50}$ = 124 μ g/ml) as well as induces apoptosis and cell cycle arrest at the G_2/M phase in the same cells.¹ Koumine (200 and 400 μ g/ml) reduces LPS-induced increases in inducible nitric oxide synthase (iNOS), IL-1β, IL-6, and TNF-α levels in RAW 264.7 macrophages.² It inhibits acetic acid-induced writhing and formalin-induced paw licking and biting in mice when administered at doses of 2 and 10 mg/kg.³ Koumine (0.4-10 mg/kg) increases the number of entries into and percentage of time spent in the open arms of the elevated plus maze in mice, indicating anxiolytic-like activity.4

References

- 1. Zhang, X., Chen, Y., Gao, B., et al. Apoptotic effect of koumine on human breast cancer cells and the mechanism involved. Cell Biochem. Biophys. 72(2), 411-416 (2015).
- 2. Yuan, Z., Matias, F.B., Wu, J., et al. Koumine attenuates lipopolysaccaride-stimulated inflammation in RAW264.7 macrophages, coincidentally associated with inhibition of NF-κB, ERK and p38 pathways. Int. J. Mol. Sci. 17(3), 430 (2016).
- 3. Liu, M., Huang, H.-H., Yang, J., et al. The active alkaloids of Gelsemium elegans Benth. are potent anxiolytics. Psychopharmacology (Berl) 225(4), 839-851 (2013).
- 4. Xu, Y., Qiu, H.-Q., Liu, H., et al. Effects of koumine, an alkaloid of Gelsemium elegans Benth., on inflammatory and neuropathic pain models and possible mechanism with allopregnanolone. Pharmacol. Biochem. Behav. 101(3), 504-514 (2012).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 12/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM