

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## 1-Arachidonoyl-d<sub>5</sub>-rac-glycerol

Item No. 362152

Formal Name: 5Z,8Z,11Z,14Z-eicosatetraenoic

acid, 1-glycerol-1,1,2,3,3-d<sub>5</sub> ester

Synonym: 1-AG-d<sub>5</sub> MF:  $C_{23}H_{33}D_5O_4$ 

FW: 383.6 **Chemical Purity:** ≥97% (1-AG)

Deuterium

Incorporation: ≥99% deuterated forms (d<sub>1</sub>-d<sub>5</sub>);

Supplied as: A solution in acetonitrile

Storage: -80°C Stability: ≥6 months

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



1-Arachidonyl- $d_s$ -rac-glycerol (1-AG- $d_s$ ) is intended for use as an internal standard for the quantification of 1-AG (Item No. 62150) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

1-AG-d<sub>5</sub> is supplied as a solution in acetonitrile. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of  $1-AG-d_5$  in these solvents is approximately 2 mg/ml.

### Description

The glyceryl esters of arachidonic acid have been proposed as endogenous cannabinoid ligands. 2-AG is 10 to 100 times more potent than 1-AG in ligand binding affinity and agonist activity at the CB₁ receptor,<sup>2</sup> and is thus considered to be the natural ligand. However, 2-AG is chemically unstable and undergoes rapid isomerization to 1-AG (synonymous with 1(3)-AG) both in vitro and in vivo. 1-AG is a frequent contaminant in synthetic 2-AG preparations, and can markedly reduce their cannabinergic potency. 1-AG is a weak CB<sub>1</sub> receptor agonist and may have other pharmacologic properties.

### References

- 1. Sugiura, T., Kodaka, T., Kondo, S., et al. 2-Arachidonoylglycerol, a putative endogenous cannabinoid receptor ligand, induces rapid, transient elevation of intracellular free Ca<sup>2+</sup> in neuroblastoma X glioma hybrid NG108-15 cells. Biochem. Biophys. Res. Commun. 229, 58-64 (1996).
- 2. Sugiura, T., Kodaka, T., Kondo, S., et al. Is the cannabinoid CB<sub>1</sub> receptor a 2-arachidonoylglycerol receptor? Structural requirements for triggering a Ca<sup>2+</sup> transient in NG108-15 cells. J. Biochem. 122, 890-895 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 05/23/2019

## **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM