

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



2-Arachidonoyl Glycerol-d₅

Item No. 362162

Formal Name: 5Z,8Z,11Z,14Z-eicosatetraenoic acid,

2-glycerol-1,1,2,3,3-d₅ ester

Synonym: 2-AG-d₅ MF: $C_{23}H_{33}D_5O_4$ FW: 383.6

Chemical Purity: ≥95% (2-Arachidonoyl Glycerol; as a

9:1 mixture of 2-AG and 1-AG)

Deuterium

 \geq 99% deuterated forms (d₁-d₅); \leq 1% d₀ Incorporation:

Supplied as: A solution in acetonitrile

Storage: -80°C Stability: ≥6 months Special Conditions: Light sensitive

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



2-Arachidonoyl glycerol- d_5 (2-AG- d_5) is intended for use as an internal standard for the quantification of 2-AG (Item No. 62160) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

2-AG-d₅ is supplied as a solution in acetonitrile. To change the solvent, simply evaporate the acetonitrile under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO purged with an inert gas can be used at a concentration of approximately 10 mg/ml. 2-AG-d₅ is also miscible in ethanol.

Description

2-AG is an endogenous agonist of the CB₁ receptor. 1,2 Unlike anandamide, 2-AG is present at relatively high levels in the central nervous system; it is the most abundant molecular species of monoacylglycerol found in rat brain. 1,3 Formation of 2-AG is calcium-dependent and is mediated by the activities of PLC and DAG lipase.¹ 2-AG acts as a full agonist at the CB₁ receptor. At a concentration of 0.3 nM, 2-AG induces a rapid, transient increase in intracellular free calcium in NG108-15 neuroblastoma X glioma cells through a CB₁ receptor-dependent mechanism.² 2-AG is metabolized in vitro by MAG lipase and fatty acid amide hydrolase, with MAG lipase likely being the principle metabolizing enzyme in vivo.⁴

References

- 1. Stella, N., Schweitzer, P., and Piomelli, D. Nature 388(6644), 773-778 (1997).
- Sugiura, T., Kodaka, T., Nakane, S., et al. J. Biol. Chem. 274(5), 2794-2801 (1999).
- Kondo, S., Kondo, H., Nakane, S., et al. FEBS Lett. 429(2), 152-156 (1998).
- 4. Dinh, T.P., Carpenter, D., Leslie, F.M., et al. Proc. Natl. Acad. Sci. USA 99(16), 10819-10824 (2002).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 07/28/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM