



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Fmoc-1-Methyl-L-histidine

Item No. 36312

CAS Registry No.: 202920-22-7
Formal Name: N-[(9H-fluoren-9-ylmethoxy) carbonyl]-1-methyl-L-histidine
Synonym: Fmoc-His(1-Me)-OH
MF: C₂₂H₂₁N₃O₄
FW: 391.4
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Fmoc-1-methyl-L-histidine is supplied as a solid. A stock solution may be made by dissolving the Fmoc-1-methyl-L-histidine in the solvent of choice, which should be purged with an inert gas. Fmoc-1-methyl-L-histidine is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of Fmoc-1-methyl-L-histidine in these solvents is approximately 10 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of Fmoc-1-methyl-L-histidine can be prepared by directly dissolving the solid in aqueous buffers. The solubility of Fmoc-1-methyl-L-histidine in PBS (pH 7.2) is approximately 0.2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Fmoc-1-methyl-L-histidine is an amino acid-containing building block.¹ It has been used in the production of ϵ -nitrogen-ligated copper centers in copper nitrite reductases.

Reference

1. Koebke, K.J., Yu, F., Van Stappen, C., *et al.* Methylated histidines alter tautomeric preferences that influence the rates of Cu nitrite reductase catalysis in designed peptides. *J. Am. Chem. Soc.* **141**(19), 7765-7775 (2019).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/02/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM