

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Methylprotodioscin

Item No. 36344

CAS Registry No.:	54522-52-0	
Formal Name:	(3β,22α,25R)-26-(β-D-glucopyranosyloxy)-	
	22-methoxyfurost-5-en-3-yl O-6-deoxy-	
	α -L-mannopyranosyl-(1 \rightarrow 2)-O- β -D-	QH OH
	glucopyranoside	но.
Synonyms:	22-O-Methylprotodioscin, NSC 698790,	
	Smilax Saponin B	
MF:	$C_{52}H_{86}O_{22}$	
FW:	1,063.2	
Purity:	≥98%	
Supplied as:	A solid	
Storage:	-20°C	
Stability:	≥4 years	
Item Origin:	Plant/Dioscorea opposita	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Methylprotodioscin is supplied as a solid. A stock solution may be made by dissolving the methylprotodioscin in the solvent of choice, which should be purged with an inert gas. Methylprotodioscin is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of methylprotodioscin in these solvents is approximately 2 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of methylprotodioscin can be prepared by directly dissolving the solid in aqueous buffers. The solubility of methylprotodioscin in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Methylprotodioscin is a steroidal saponin that has been found in A. cochinchinensis and has diverse biological activities.¹⁻³ It suppresses mucin 5AC (MUC5AC) production induced by EGF or phorbol 12-myristate 13-acetate (PMA; Item No. 10008014) in H292 cells when used at a concentration of 1 μ M.¹ Methylprotodioscin inhibits the proliferation of HeLa cervical cancer cells (IC₅₀ = 18.31 μ M) but has no effect on non-cancerous HEK293 cells.² It also induces production of reactive oxygen species (ROS), apoptosis, and cell cycle arrest at the G_2/M phase in HeLa cells. In vivo, methylprotodioscin (50 mg/kg) decreases bone loss and prevents decreases in bone mineral content and bone mineral density in a mouse model of ovariectomy-induced osteoporosis.³

References

- 1. Lee, H.J., Park, J.S., Yoon, Y.P., et al. Phytomedicine 22(5), 568-572 (2015).
- 2. Ma, Y.-L., Zhang, Y.-S., Zhang, F., et al. Food Chem. Toxicol. 132, 110655 (2019).
- 3. Yin, J., Tezuka, Y., Kouda, K., et al. Planta Med. 70(3), 220-226 (2004).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/15/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM