



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



8(S)-HETrE

Item No. 36360

CAS Registry No.: 889573-69-7
Formal Name: 8S-hydroxy-9E,11Z,14Z-eicosatrienoic acid
MF: C₂₀H₃₄O₃
FW: 322.5
Purity: ≥98%
UV/Vis.: λ_{max}: 236 nm ε: 23,000
Supplied as: A solution in ethanol
Storage: -20°C
Stability: ≥1 year
Special Conditions: Oxygen and light sensitive



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

8(S)-HETrE is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 8(S)-HETrE in these solvents is miscible.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 8(S)-HETrE is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 8(S)-HETrE in PBS, pH 7.2, is approximately 0.8 mg/ml. For greater aqueous solubility, 8(S)-HETrE can be directly dissolved in 0.1 M Na₂CO₃ (solubility of 2 mg/ml) and then diluted with PBS (pH 7.2) to achieve the desired concentration or pH. We do not recommend storing the aqueous solution for more than one day.

Description

8(S)-HETrE is a monohydroxy polyunsaturated fatty acid produced by rabbit neutrophil lipoxygenase when dihomo-γ-linolenic acid (DGLA; Item No. 90230) is used as a substrate.¹ Although the biological activities of 8(S)-HETrE have not been well characterized, it is expected to behave similarly to 8(S)-HETE (Item No. 34360).

Reference

1. Borgeat, P., Hamberg, M., and Samuelsson, B. Transformation of arachidonic acid and homo-γ-linolenic acid by rabbit polymorphonuclear leukocytes. Monohydroxy acids from novel lipoxygenases. *J. Biol. Chem.* **251**(24), 7816-7820 (1976).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/02/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM