

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## Phenylacetylglycine

Item No. 36368

CAS Registry No.: 500-98-1

Formal Name: N-(2-phenylacetyl)-glycine Synonyms: NSC 408424, NSC 92778,

PAG, PAGly, Phenaceturic Acid,

N-Phenylacetylglycine

MF:  $C_{10}H_{11}NO_3$ FW: 193.2 **Purity:** ≥98% Supplied as: A solid Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### **Laboratory Procedures**

Phenylacetylglycine is supplied as a solid. A stock solution may be made by dissolving the phenylacetylglycine in the solvent of choice, which should be purged with an inert gas. Phenylacetylglycine is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of phenylacetylglycine in these solvents is approximately 1, 10, and 25 mg/ml, respectively.

#### Description

Phenylacetylglycine is a glycine conjugate of phenylacetate. 1-4 It is formed via the conversion of phenylalanine by gut microbiota to phenylacetic acid followed by glycine conjugation in the liver.<sup>4,5</sup> Phenylacetylglycine (100 μM) decreases hypoxia-induced apoptosis in neonatal mouse cardiomyocytes.<sup>3</sup> Levels of phenylacetylglycine in urine and plasma are increased in a rat model of phospholipidosis induced by the antiarrhythmic agent amiodarone (Item No. 15213). Phenylacetylglycine reduces myocardial infarct size in a mouse model of ischemia-reperfusion injury.<sup>3</sup>

#### References

- 1. Kamiguchi, H., Murabayashi, M., Mori, I., et al. Biomarker discovery for drug-induced phospholipidosis: Phenylacetylglycine to hippuric acid ratio in urine and plasma as potential markers. Biomarkers 22(2), 178-188 (2016).
- 2. Jones, A.R. Some observations on the urinary excretion of glycine conjugates by laboratory animals. Xenobiotica 12(6), 387-395 (1982).
- 3. Xu, X., Lu, W.-J., Shi, J.-Y., et al. The gut microbial metabolite phenylacetylglycine protects against cardiac injury caused by ischemia/reperfusion through activating β2AR. Arch. Biochem. Biophys. 697, 108720 (2021).
- 4. Nemet, I., Saha, P.P., Gupta, N., et al. A cardiovascular disease-linked gut microbial metabolite acts via adrenergic receptors. Cell 180(5), 862-877.e22 (2020).
- Zhu, Y., Dwidar, M., Nemet, I., et al. Two distinct gut microbial pathways contribute to meta-organismal production of phenylacetylglutamine with links to cardiovascular disease. Cell Host Microbe 31(1), 18-32.e9 (31(1)).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 04/18/2023

## **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM