



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



2-Methylbutyrylglycine

Item No. 36379

CAS Registry No.: 52320-67-9
Formal Name: N-(2-methyl-1-oxobutyl)-glycine
Synonyms: 2-MBG, N-sec-Valerylglycine
MF: C₇H₁₃NO₃
FW: 159.2
Purity: ≥95%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

2-Methylbutyrylglycine is supplied as a solid. A stock solution may be made by dissolving the 2-methylbutyrylglycine in the solvent of choice, which should be purged with an inert gas. 2-Methylbutyrylglycine is soluble in organic solvents such as ethanol and DMSO.

Description

2-Methylbutyrylglycine is a metabolite of the essential amino acid L-isoleucine.^{1,2} It is produced by hydrolysis of the L-isoleucine catabolic intermediate (S)-2-methylbutyryl-CoA, which accumulates when the activity of 2-methylbutyryl-CoA dehydrogenase is deficient. 2-Methylbutyrylglycine (0.5-5 mM) induces the formation of thiobarbituric acid reactive substances (TBARS) in isolated rat cerebral cortex.³ Elevated levels of 2-methylbutyrylglycine are associated with 2-methylbutyryl-CoA dehydrogenase deficiency (2-MBCDD), also known as short/branched-chain acyl-CoA dehydrogenase (SBCAD) deficiency.^{1,2}

References

1. Gibson, K.M., Burlingame, T.G., Hogema, B., *et al.* 2-Methylbutyryl-coenzyme A dehydrogenase deficiency: A new inborn error of L-isoleucine metabolism. *Pediatr. Res.* **47(6)**, 830-833 (2000).
2. Fong, B.M.-W., Tam, S., and Leung, S.-Y. Quantification of acylglycines in human urine by HPLC electrospray ionization-tandem mass spectrometry and the establishment of pediatric reference interval in local Chinese. *Talanta* **88**, 193-200 (2012).
3. Knebel, L.A., Zanatta, Á., Tonin, A.M., *et al.* 2-Methylbutyrylglycine induces lipid oxidative damage and decreases the antioxidant defenses in rat brain. *Brain Res.* **1478**, 74-82 (2012).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/06/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM