

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



KAS 08

Item No. 36474

CAS Registry No.: 203861-20-5

Formal Name: (4S)-1-[[2,3-dihydro-1-(4-nitrobenzoyl)-

1H-indol-5-yl]sulfonyl]-4-phenyl-2-

imidazolidinone

MF: $C_{24}H_{20}N_4O_6S$

492.5 FW: ≥95% **Purity:** UV/Vis.: λ_{max} : 272 nm Supplied as: A solid Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

KAS 08 is supplied as a solid. A stock solution may be made by dissolving the KAS 08 in the solvent of choice, which should be purged with an inert gas. KAS 08 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of KAS 08 in ethanol and DMSO is approximately 12 mg/ml and approximately 16 mg/ml in DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of KAS 08 can be prepared by directly dissolving the solid in aqueous buffers. The solubility of KAS 08 in PBS (pH 7.2) is approximately 0.25. We do not recommend storing the aqueous solution for more than one day.

Description

KAS 08 is an activator of stimulator of interferon genes (STING).¹ It activates STING with an EC₅₀ value of 0.33 μM in a reporter assay using THP-1 human monocytes. KAS 08 increases the levels of IFN-β and IP-10 in the presence, but not absence, of cGAMP. It reduces the viability of CT26 mouse colorectal cancer cells when used at concentrations ranging from 0.31 to 20 μM. KAS 08 (15 mg/kg), when administered in combination with exogenous 2'3'-cGAMP (Item No. 19887), reduces tumor growth in a CT26 mouse syngeneic model.

Reference

1. Jung, H.R., Jo, S., Jeon, M.J., et al. Development of small-molecule STING activators for cancer immunotherapy. Biomedicines 10(1), 33 (2022).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/17/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM