



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



3,5,6,7,8,3',4'-Heptamethoxyflavone

Item No. 36475

CAS Registry No.:	1178-24-1
Formal Name:	2-(3,4-dimethoxyphenyl)-3,5,6,7,8-pentamethoxy-H-1-benzopyran-4-one
Synonyms:	HMF, NSC 618928
MF:	C ₂₂ H ₂₄ O ₉
FW:	432.4
Purity:	≥98%
Supplied as:	A solid
Storage:	-20°C
Stability:	≥4 years
Item Origin:	Plant/Citrus sp.



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

3,5,6,7,8,3',4'-Heptamethoxyflavone (HMF) is supplied as a solid. A stock solution may be made by dissolving the HMF in the solvent of choice, which should be purged with an inert gas. HMF is soluble in methanol and DMSO.

Description

HMF is a polymethoxyflavone that has been found in *C. reticulata* and has diverse biological activities.¹⁻⁴ It inhibits the efflux of Hoechst 33342 (Item No. 15547) by breast cancer resistance protein (BCRP) and rhodamine 123 (Item No. 16672) by P-glycoprotein (P-gp), also known as multidrug resistance protein 1 (MDR1), in MDCK cells expressing the human transporters (IC₅₀s = 1.4 and 31 μM, respectively).¹ HMF (50, 100, and 200 μg/ml) prevents UVB-induced decreases in type I procollagen in primary human dermal fibroblast neonatal (HDFn) cells.² It inhibits the proliferation of B16 melanoma 4A5, CCRF-HSB-2 leukemia, and TGBC11TKB gastric cancer cells (IC₅₀s = 38, 29, and 9 μM, respectively).³ HMF (50 mg/kg) prevents corticosterone-induced increases in immobility time in the forced swim test in mice.⁴

References

1. Pick, A., Müller, H., Mayer, R., et al. Structure-activity relationships of flavonoids as inhibitors of breast cancer resistance protein (BCRP). *Bioorg. Med. Chem.* **19(6)**, 2090-2102 (2011).
2. Kim, H.-I., Jeong, Y.-U., Kim, J.-H., et al. 3,5,6,7,8,3',4'-Heptamethoxyflavone, a *Citrus* flavonoid, inhibits collagenase activity and induces type I procollagen synthesis in HDFn cells. *Int. J. Mol. Sci.* **19(2)**, 620 (2018).
3. Kawaii, S., Tomono, Y., Katase, E., et al. Antiproliferative activity of flavonoids on several cancer cell lines. *Biosci. Biotechnol. Biochem.* **63(5)**, 896-899 (1999).
4. Sawamoto, A., Okuyama, S., Yamamoto, K., et al. 3,5,6,7,8,3',4'-Heptamethoxyflavone, a citrus flavonoid, ameliorates corticosterone-induced depression-like behavior and restores brain-derived neurotrophic factor expression, neurogenesis, and neuroplasticity in the hippocampus. *Molecules* **21(4)**, 541 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/17/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM