



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Ciraparantag (trifluoroacetate salt)

Item No. 36809

Formal Name: (2S,2'S)-N,N'-(1,4-piperazinediyldi-3,1-propanediyl)bis[2-amino-5-[(aminoiminomethyl)amino]pentanamide, trifluoroacetate salt
Synonym: PER977
MF: C₂₂H₄₈N₁₂O₂ • XCF₃COOH
FW: 512.7
Purity: ≥95%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Ciraparantag (trifluoroacetate salt) is supplied as a solid. A stock solution may be made by dissolving the ciraparantag (trifluoroacetate salt) in the solvent of choice, which should be purged with an inert gas. It is also soluble in water. The solubility of ciraparantag (trifluoroacetate salt) in methanol and water is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Ciraparantag is an anticoagulant reversal agent.^{1,2} It selectively binds to unfractionated heparin, enoxaparin, and Factor IXa (K_ds = 28, 17, and 35 μM, respectively) over edoxaban (Item No. 24030), rivaroxaban (Item No. 16043), Factor Xa, Factor IX, thrombin (Item No. 13188), antithrombin, and fibrinogen, and only weakly binds to an enoxaparin-antithrombin complex and fondaparinux (Item No. 24033; K_ds = 690 and 9,600 μM, respectively), in cell-free assays.¹ Ciraparantag (200 μg/ml) prevents decreases in fibrin diameter induced by the Factor Xa inhibitor edoxaban in isolated human whole blood. It reduces edoxaban-, dabigatran-, apixaban-, rivaroxaban-, unfractionated heparin-, or enoxaparin-stimulated increases in tail transection-induced blood loss in rats.²

References

1. Kalathottukaren, M.T., Creagh, A.L., Abbina, S., *et al.* Comparison of reversal activity and mechanism of action of UHRA, andexanet, and PER977 on heparin and oral FXa inhibitors. *Blood Adv.* **2(16)**, 2104-2114 (2018).
2. Ansell, J., Laulicht, B.E., Bakhru, S.H., *et al.* Ciraparantag, an anticoagulant reversal drug: Mechanism of action, pharmacokinetics, and reversal of anticoagulants. *Blood* **137(1)**, 115-125 (2021).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/27/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM