

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Blarcamesine

Item No. 36812

CAS Registry No.:		I
Formal Name:	tetrahydro-N,N-dimethyl-2,2-diphenyl-3-	
	furanmethanamine	
Synonyms:	AE37, AV2-73	
MF:	C ₁₉ H ₂₃ NO	\leq
FW:	281.4	\sim
Purity:	≥98%	
Supplied as:	A crystalline solid	0/
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis		

Laboratory Procedures

Blarcamesine is supplied as a crystalline solid. A stock solution may be made by dissolving the blarcamesine in the solvent of choice, which should be purged with an inert gas. Blarcamesine is soluble in organic solvents such as ethanol and dimethyl formamide. The solubility of blarcamesine in these solvents is approximately 30 mg/ml. Blarcamesine is slightly soluble in DMSO.

Description

Blarcamesine is an agonist of the sigma non-opioid intracellular receptor 1 (σ_1 receptor).¹ It inhibits reductions in mitochondrial oxygen consumption and lipid peroxidation induced by amyloid- β (A β) (25-35) (Aβ (25-35)) in isolated mouse hippocampus when administered at a dose of 1 mg/kg. Blarcamesine (1 mg/kg) improves deficits in novel object recognition induced by A β (25-35) in mice, as well as reduces tau phosphorylation and A β (1-42) levels in the hippocampus in the same model.² It increases the latency to fall in the rotarod test, reduces gait impairments, and reduces increases in the mean acoustic startle response in an Mecp2^{-/-} mouse model of Rett syndrome, an X-linked neurodevelopmental disorder, when administered at a dose of 30 mg/kg per day.³

References

- 1. Lahmy, V., Long, R.M., Morin, D., et al. Mitochondrial protection by the mixed muscarinic/ σ_1 ligand ANAVEX2-73, a tetrahydrofuran derivative, in $A\beta_{25-35}$ peptide-injected mice, a nontransgenic Alzheimer's disease model. Front. Cell. Neurosci. 8, 463 (2015).
- 2. Lahmy, V., Meunier, J., Malström, S., et al. Blockade of Tau hyperphosphorylation and Aβ₁₋₄₂ generation by the aminotetrahydrofuran derivative ANAVEX2-73, a mixed muscarinic and σ_1 receptor agonist, in a nontransgenic mouse model of Alzheimer's disease. Neuropsychopharmacology 38(9), 17061723 (2013).
- 3. Kaufmann, W., Sprouse, J., Rebowe, N., et al. ANAVEX[®]2-73 (blarcamesine), a Sigma-1 receptor agonist, ameliorates neurologic impairments in a mouse model of Rett syndrome. Pharmacol. Biochem. Behav. 187, 172796 (2019).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/11/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM