

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



NG-497

Item No. 36886

CAS Registry No.: 2598242-66-9

Formal Name: 6-(4-ethoxyphenyl)-4-

methoxy-2-pyridinecarboxylic


acid, 1-methylethyl ester

MF: C<sub>18</sub>H<sub>21</sub>NO<sub>4</sub> 315.4 FW: **Purity:** ≥98%

UV/Vis.:  $\lambda_{\text{max}}$ : 256, 282 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



### **Laboratory Procedures**

NG-497 is supplied as a crystalline solid. A stock solution may be made by dissolving the NG-497 in the solvent of choice, which should be purged with an inert gas. NG-497 is soluble in the organic solvent DMSO at a concentration of approximately 3 mg/ml.

#### Description

NG-497 is an inhibitor of adipose triglyceride lipase (ATGL; IC $_{50}$  = 1  $\mu$ M), also known as patatin-like phospholipase domain-containing protein 2 (PNPLA2).1 It is selective for ATGL over PNPLA1, PNPLA3, PNPLA4, and PNPLA5 at 100 μM. NG-497 selectively inhibits the triacylglycerol hydrolase activity of human and rhesus monkey ATGL over rat, dog, and marmoset ATGL, which it inhibits by less than 20% at 50 μM, and over mouse, goat, and pig ATGL at which it is inactive at 50 µM. NG-497 inhibits fatty acid and glycerol release induced by isoproterenol (Item No. 15592) in human Simpson-Golabi-Behmel syndrome (SGBS) adipocytes (IC $_{50}$  = 1.5  $\mu$ M for both). It also inhibits forskolin-induced fatty acid release in primary human adipocytes.

#### Reference

1. Grabner, G.F., Guttenberger, N., Mayer, N., et al. Small-molecule inhibitors targeting lipolysis in human adipocytes. J. Am. Chem. Soc. 144(14), 6237-6250 (2022).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 08/12/2022

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM