



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## Piperazine Erastin

Item No. 36934

**CAS Registry No.:** 1538593-71-3  
**Formal Name:** 2-[[4-[2-(4-chlorophenoxy)acetyl]-1-piperazinyl]methyl]-3-[2-(1-methylethoxy)-5-(1-piperazinylmethyl)phenyl]-4(3H)-quinazolinone

**MF:** C<sub>35</sub>H<sub>41</sub>ClN<sub>6</sub>O<sub>4</sub>

**FW:** 645.2


**Purity:** ≥98%

**UV/Vis.:** λ<sub>max</sub>: 227 nm

**Supplied as:** A solid

**Storage:** -20°C

**Stability:** ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Piperazine erastin is supplied as a solid. A stock solution may be made by dissolving the piperazine erastin in the solvent of choice, which should be purged with an inert gas. Piperazine erastin is soluble in the organic solvent DMSO at a concentration of approximately 10 mg/ml.

### Description

Piperazine erastin is an inducer of ferroptosis.<sup>1,2</sup> It inhibits glutamate release in human CCF-STTG1 astrocytoma cells (IC<sub>50</sub> = 0.8 μM), indicating inhibition of the system x<sub>c</sub><sup>-</sup> cystine/glutamate transporter, and decreases glutathione (GSH) levels in HT-1080 fibrosarcoma cells when used at a concentration of 10 μM. Piperazine erastin inhibits the growth of BJeLR cells overexpressing the H-Ras activating mutation H-Ras<sup>G12V</sup> (IC<sub>50</sub> = 0.3 μM).<sup>1</sup> It increases hepatic mRNA expression of the gene encoding Cox-2 and prevents tumor formation in an HT-1080 mouse xenograft model when administered at a dose of 60 mg/kg.<sup>2</sup>

### References

1. Larraufie, M.-H., Yang, W.S., Jiang, E., *et al.* Incorporation of metabolically stable ketones into a small molecule probe to increase potency and water solubility. *Bioorg. Med. Chem. Lett.* **25(21)**, 4787-4792 (2015).
2. Yang, W.S., SriRamaratnam, R., Welsch, M.E., *et al.* Regulation of ferroptotic cancer cell death by GPX4. *Cell* **156(1-2)**, 317-331 (2014).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/21/2022

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM