



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Layilin Isoform 2 Extracellular Domain (human, recombinant)


Item No. 37021

Overview and Properties

Synonym:	LAYN
Source:	Active recombinant human C-terminal His-tagged coagulation factor IX expressed in HEK293 cells
Amino Acids:	22-220
Uniprot No.:	Q6UX15
Molecular Weight:	49.5 kDa
Storage:	-80°C (as supplied)
Stability:	≥1 year
Purity:	≥90% estimated by SDS-PAGE
Supplied in:	Lyophilized from sterile 50 mM Tris, pH 7.5, with 100 mM glycine and 10 mM sodium chloride
Endotoxin Testing:	<1.0 EU/μg, determined by the LAL endotoxin assay
Bioactivity:	See figures for details

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Image



Lane 1: MW Markers

Lane 2: Layilin Isoform 2 Extracellular Domain

SDS-PAGE Analysis of Layilin Isoform 2 Extracellular Domain.

This protein has a calculated molecular weight of 49.5 kDa. It has an apparent molecular weight of approximately 60-65 kDa by SDS-PAGE under reducing conditions due to glycosylation.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/27/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Layilin is a C-type lectin receptor.¹ It is composed of an extracellular region that contains the C-type lectin homology domain, a transmembrane domain, and a cytoplasmic domain. Layilin is ubiquitously expressed and localizes to the cell surface, but has also been found in mitochondria.^{1,2} It is involved in cell migration, the epithelial-to-mesenchymal transition (EMT), and antitumor immunity and binds to the extracellular matrix proteins hyaluronan and glycosylated collagen, as well as the cytoskeleton-membrane linker proteins talin, merlin, and radixin.^{1,3-5} The layilin isoform 2 extracellular domain binds to melanoma cells in a calcium-dependent manner.⁵ Knockout of *LAYN* prevents the accumulation of lymph node-derived CD8⁺ tumor-infiltrating lymphocytes (TILs) in a competitive adoptive transfer murine colon adenocarcinoma model.⁶ Layilin protein levels are increased in CD8⁺ TILs that highly express programmed cell death protein 1 (PD-1) and cytotoxic T lymphocyte protein 4 (CTLA-4) isolated from patients with metastatic melanoma. Cayman's Layilin Isoform 2 Extracellular Domain (human, recombinant) protein consists of 437 amino acids, has a calculated molecular weight of 49.5 kDa, and a predicted N-terminus of Ala22 after signal peptide cleavage. By SDS-PAGE, under reducing conditions, the apparent molecular mass of the protein is 60-65 kDa due to glycosylation.

References

1. Borowsky, M.L. and Hynes, R.O. Layilin, a novel talin-binding transmembrane protein homologous with C-type lectins, is localized in membrane ruffles. *J. Cell Biol.* **143**(2), 429-442 (1998).
2. Tsutiya, A., Arito, M., Tagashira, T., *et al.* Layilin promotes mitochondrial fission by cyclin-dependent kinase 1 and dynamin-related protein 1 activation in HEK293T cells. *Biochem. Biophys. Res. Commun.* **549**, 143-149 (2021).
3. Adachi, T., Arito, M., Suematsu, N., *et al.* Roles of layilin in TNF- α -induced epithelial-mesenchymal transformation of renal tubular epithelial cells. *Biochem. Biophys. Res. Commun.* **467**(1), 63-69 (2015).
4. Carvalho, A.M., Reis, R.L., and Pashkuleva, I. Hyaluronan receptors as mediators and modulators of the tumor microenvironment. *Adv. Healthc. Mater.* e2202118 (2022).
5. Glasgow, J.E., Byrnes, J.R., Barbee, S.D., *et al.* Identifying and antagonizing the interactions between layilin and glycosylated collagens. *Cell Chem. Biol.* **29**(4), 597-604 (2022).
6. Mahuron, K.M., Moreau, J.M., Glasgow, J.E., *et al.* Layilin augments integrin activation to promote antitumor immunity. *J. Exp. Med.* **217**(9), e20192080 (2020).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM