

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


PRODUCT INFORMATION



N¹-Acetylputrescine

Item No. 37185

| CAS Registry No.: Formal Name: | 5699-41-2 N-(4-aminobutyl)-acetamide |
|--|--|
| Synonyms: | Monoacetylputrescine, N-acetyl Putrescine |
| MF: | $C_6H_{14}N_{20}$ |
| FW: | 130.2 N/ / / / / / / / / / / / / / / / / / / |
| Purity: | ≥95% |
| Supplied as: | A neat liquid |
| Storage: | -20°C |
| Stability: | ≥4 years |
| Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis. | |



Laboratory Procedures

N¹-Acetylputrescine is supplied as a neat liquid. A stock solution may be made by dissolving the N^1 -acetylputrescine in the solvent of choice, which should be purged with an inert gas. N^1 -Acetylputrescine is soluble in organic solvents such as ethanol and DMSO. The solubility of N¹-acetylputrescine in these solvents is approximately 16 and 10 mg/ml, respectively. N¹-Acetylputrescine is also slightly soluble in dimethyl formamide.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of N¹-acetylputrescine can be prepared by directly dissolving the neat liquid in aqueous buffers. The solubility of N^1 -acetylputrescine in PBS (pH 7.2) is approximately 3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

N¹-Acetylputrescine is a metabolite of the endogenous polyamine putrescine.¹ It is formed from putrescine by diamine acetyltransferase 1 in the brain. Serum levels of N¹-acetylputrescine are increased in patients with breast cancer.²

References

- 1. Seiler, N. and Al-Therib, M.J. Putrescine catabolism in mammalian brain. Biochem. J. 144(1), 29-35 (1974).
- 2. Lee, Y.R., Lee, J.W., Hong, J., et al. Simultaneous determination of polyamines and steroids in human serum from breast cancer patients using liquid chromatography-tandem mass spectrometry. Molecules 26(4), 1153 (2021).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/17/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM