

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Sulfadoxin-d₄

Item No. 37200

CAS Registry No.:	1330266-05-1	
Formal Name:	4-amino-N-(5,6-dimethoxy-4-pyrimidinyl)-	
	benzenesulfonamide-d ₄	
Synonym:	Ro 4-4393-d ₄	
MF:	$C_{12}H_{10}D_4N_4O_4S$	
FW:	314.4	
Chemical Purity:	≥98% (Sulfadoxin)	
Deuterium		
Incorporation:	≥99% deuterated forms (d ₁ -d ₄); ≤1% d ₀	H_2N^2
Supplied as:	A solid	Ď
Storage:	-20°C	
Stability:	≥4 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Sulfadoxin- d_{4} is intended for use as an internal standard for the quantification of sulfadoxin (Item No. 21488) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Sulfadoxin- d_4 is supplied as a solid. A stock solution may be made by dissolving the sulfadoxin- d_4 in the solvent of choice, which should be purged with an inert gas. Sulfadoxin- d_4 is soluble in DMSO, methanol, and acetone.

Description

Sulfadoxin is a sulfonamide antibiotic that inhibits dihydropteroate synthetase (DHPS), an enzyme involved in folic acid (Item No. 20515) synthesis.¹ Sulfadoxin competes with 4-aminobenzoate (PABA), the native substrate of DHPS, and inhibits PABA incorporation into folic acid.² Folate is essential for purine and pyrimidine synthesis, therefore, sulfadoxin has antiproliferative activity in non-resistant P. falciparum.^{2,3} Sulfadoxin inhibits growth of P. falciparum in vitro, but potency varies between non-resistant (IC_{50} = 4 ng/ml) and resistant strains (IC₅₀ = 3,970 ng/ml).³

References

- 1. Brooks, D.R., Wang, P., Read, M., et al. Sequence variation of the hydroxymethyldihydropterin pyrophosphokinase: dihydropteroate synthase gene in lines of the human malaria parasite, Plasmodium falciparum, with differing resistance to sulfadoxine. Eur. J. Biochem. 224(2), 397-405 (1994).
- Hyde, J.E. Exploring the folate pathway in Plasmodium falciparum. Acta. Trop. 94(3), 191-206 (2005).
- 3. Wang, P., Read, M., Sims, P.F.G., et al. Sulfadoxine resistance in the human malaria parasite Plasmodium falciparum is determined by mutations in dihydropteroate synthetase and an additional factor associated with folate utilization. Mol. Microbiol. 23(5), 979-986 (1997).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/21/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM