



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# PRODUCT INFORMATION

## (±)-10-hydroxy-12(Z)-Octadecenoic Acid

Item No. 37281

CAS Registry No.: 34932-12-2

Formal Name: 10-hydroxy-12Z-octadecenoic acid

Synonym: 10-hydroxy-cis-12-Octadecenoic Acid

MF: C<sub>18</sub>H<sub>34</sub>O<sub>3</sub>

FW: 298.5

Purity: ≥98%

Supplied as: A solution in ethanol

Storage: -20°C

Stability: ≥1 year



*Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.*

### Description

(±)-10-hydroxy-12(Z)-Octadecenoic acid is an oxylipin and metabolite of linoleic acid (Item Nos. 90150 | 90150.1 | 21909).<sup>1</sup> It is formed from linoleic acid by conjugated linoleic acid-hydrolase (CLA-HY) as an intermediate in CLA biosynthesis and can also be produced from linoleic acid by gut microbiota. (±)-10-hydroxy-12(Z)-Octadecenoic acid (30 µM) inhibits LPS-induced nitric oxide (NO) production, ERK phosphorylation, and increases in inducible NO synthase (iNOS) levels in BV-2 microglia cells.<sup>2</sup> It reduces TNF-α, NO<sub>2</sub>, and IL-10 levels in LPS-stimulated and -unstimulated isolated dendritic bone marrow cells when used at a concentration of 100 µM.<sup>3</sup> (±)-10-hydroxy-12(Z)-Octadecenoic acid (100 µM) decreases LPS-induced maturation of dendritic cells in isolated mouse bone marrow cells.

### References

1. Kishino, S., Takeuchi, M., Park, S.-B., et al. Polyunsaturated fatty acid saturation by gut lactic acid bacteria affecting host lipid composition. *Proc. Natl. Acad. Sci. USA* **110**(44), 17808-17813 (2013).
2. Ikeguchi, S., Izumi, Y., Kitamura, N., et al. Inhibitory effect of the gut microbial linoleic acid metabolites, 10-oxo-trans-11-octadecenoic acid and 10-hydroxy-cis-12-octadecenoic acid, on BV-2 microglial cell activation. *J. Pharmacol. Sci.* **138**(1), 9-15 (2018).
3. Bergamo, P., Luongo, D., Miyamoto, J., et al. Immunomodulatory activity of a gut microbial metabolite of dietary linoleic acid, 10-hydroxy-cis-12-octadecenoic acid, associated with improved antioxidant/detoxifying defences. *J. Funct. Foods* **11**, 192-202 (2014).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 - USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM