



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Bradykinin (human, mouse, rat, bovine) (acetate)

Item No. 37408

CAS Registry No.: 6846-03-3

Synonyms: Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg-OH,
BK, RPPGFSPFR-OH

MF: $C_{50}H_{73}N_{15}O_{11} \cdot XC_2H_4O_2$

FW: 1,060.2

H—Arg—Pro—Pro—Gly—Phe—Ser—Pro—Phe—Arg—OH

Purity: >98%

• XCH_3CO_2H

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Bradykinin (human, mouse, rat, bovine) (acetate) is supplied as a solid. A stock solution may be made by dissolving the bradykinin (human, mouse, rat, bovine) (acetate) in the solvent of choice, which should be purged with an inert gas. Bradykinin (human, mouse, rat, bovine) (acetate) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of bradykinin (human, mouse, rat, bovine) (acetate) in these solvents is approximately 10, 12, and 5 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of bradykinin (human, mouse, rat, bovine) (acetate) can be prepared by directly dissolving the solid in aqueous buffers. The solubility of bradykinin (human, mouse, rat, bovine) (acetate) in PBS (pH 7.2) is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Bradykinin is an endogenous vasodilator that is also involved in inflammation.^{1,2} It is formed from kininogen via cleavage of high molecular weight kininogen (HMWK) by kallikrein or prekallikrein in a Factor XIIa-dependent or -independent manner.³ Bradykinin binds to the bradykinin B₂ receptor (IC₅₀ = 0.54 nM in COS-7 cells expressing the human receptor) and acts as an agonist but does not bind the bradykinin B₁ receptor (K_i = >10,000 nM in HEK293 cells expressing the human receptor).^{4,5} It stimulates arachidonic acid (Item Nos. 90010 | 90010.1 | 10006607) release in CHO cells expressing the human bradykinin B₂ receptor (EC₅₀ = 0.7 nM).⁴ Bradykinin (100 nM) induces relaxation of isolated porcine coronary arterial rings precontracted with acetylcholine (ACh; Item No. 23829) or phorbol 12,13-dibutyrate (PDBu; Item No. 27785).¹ It induces hyperalgesia in mice when administered at a dose of 1 µg/paw, an effect that can be reversed by the bradykinin B₂ antagonist icatibant (HOE 140; Item No. 24083).² Plasma levels of bradykinin are increased in patients during an acute attack of angioedema.⁶

References

1. Weintraub, N.L., Fang, X., Kaduce, T.L., et al. *Circ. Res.* **81**, 258-267 (1997).
2. Ferreira, S.H., Lorenzetti, B.B., and Poole, S. *Br. J. Pharmacol.* **110(3)**, 1227-1231 (1993).
3. Joseph, K., Tholanikunnel, B.G., and Kaplan, A.P. *J. Allergy Clin. Immunol.* **124(1)**, 143-149 (2009).
4. Hess, J.F., Borkowski, J.A., Macneil, T., et al. *Mol. Pharmacol.* **45(1)**, 1-8 (1994).
5. Bastian, S., Loillier, B., Paquet, J.L., et al. *Br. J. Pharmacol.* **122(2)**, 393-399 (1997).
6. Nussberger, J., Cugno, M., Amstutz, C., et al. *Lancet* **351(9117)**, 1693-1697 (1998).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/19/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM