

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Creatine Phosphate (sodium salt hydrate)

Item No. 37803

CAS Registry No.:	19333-65-4	
Formal Name:	N-[imino(phosphonoamino)methyl]-N-	
	methyl-glycine, disodium salt, hydrate	O NH
Synonym:	Phosphocreatine	P. L A JOH
MF:	$C_4H_{10}N_3O_5P \bullet 2Na [XH_2O]$	
FW:	255.1	Ó-
Purity:	≥95%	
Supplied as:	A solid	• 2Na ⁺ [XH ₂ O]
Storage:	-20°C	
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

Laboratory Procedures

Creatine phosphate (sodium salt hydrate) is supplied as a solid. Aqueous solutions of creatine phosphate (sodium salt hydrate) can be prepared by directly dissolving the solid in aqueous buffers. The solubility of creatine phosphate (sodium salt hydrate) in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Creatine phosphate is a phosphorylated form of creatine that is involved in energy storage in skeletal muscle and brain.^{1,2} It is formed from creatine by creatine kinase. Creatine phosphate (10 mM) reduces hydrogen peroxide- and ferrous ion-induced increases in malondialdehyde (MDA) levels in rat heart microsomes.³ It also prevents hydrogen peroxide-induced decreases in aortic flow, minute work, and peak aortic pressure in isolated perfused rat hearts when used at a concentration of 10 mM. Brain levels of creatine phosphate decrease following mild traumatic brain injury in rats.⁴

References

- 1. Wallimann, T., Wyss, M., Brdiczka, D., et al. Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: The 'phosphocreatine circuit' for cellular energy homeostasis. Biochem. J. 281(Pt 1), 21-40 (1992).
- 2. Wyss, M. and Kaddurah-Daouk, R. Creatine and creatinine metabolism. Physiol. Rev. 80(3), 1107-1213 (2000).
- 3. Zucchi, R., Poddighe, R., Limbruno, U., et al. Protection of isolated rat heart from oxidative stress by exogenous creatine phosphate. J. Mol. Cell. Cardiol. 21(1), 67-73 (1989).
- 4. Signoretti, S., Di Pietro, V., Vagnozzi, R., et al. Transient alterations of creatine, creatine phosphate, N-acetylaspartate and high-energy phosphates after mild traumatic brain injury in the rat. Mol. Cell. Biochem. 333(1-2), 269-277 (2010).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/14/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM