

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



NBQX

Item No. 38281

CAS Registry No.: 118876-58-7

Formal Name: 1,2,3,4-tetrahydro-6-nitro-2,3-dioxo-

benzo[f]quinoxaline-7-sulfonamide

Synonym: FG-9202 MF: $C_{12}H_8N_4O_6S$ FW: 336.3

Purity: ≥98%

 λ_{max} : 216, 257, 286, 380 nm UV/Vis.:

Supplied as: A solid Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Н

Laboratory Procedures

NBQX is supplied as a solid. A stock solution may be made by dissolving the NBQX in the solvent of choice, which should be purged with an inert gas. NBQX is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of NBQX in DMSO is approximately 20 mg/ml. NBQX is slightly soluble in ethanol and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of NBQX can be prepared by directly dissolving the solid in aqueous buffers. The solubility of NBQX in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

NBQX is an antagonist of AMPA and kainate receptors (IC₅₀s = 0.15 and 4.8 μ M, respectively).¹ It is selective for these receptors over NMDA receptors ($IC_{50} = \ge 90 \, \mu M$). In vivo, NBQX (20 mg/kg) increases the latency to seizures, decreases the duration of seizures, and reduces seizure severity in a rat model of epilepsy induced by pentylenetetrazole (PTZ; Item No. 18682).² It also suppresses relapse phase nicotine self-administration in rats.3

References

- 1. Sheardown, M.J., Nielsen, E.O., Hansen, A.J., et al. 2,3-Dihydroxy-6-nitro-7-sulfamoyl-benzo(F) quinoxaline: A neuroprotectant for cerebral ischemia. Science 247(4942), 571-574 (1990).
- Chen, W., Li, Y.-S., Gao, J., et al. AMPA receptor antagonist NBQX decreased seizures by normalization of perineuronal nets. PLoS One 11(11), e0166672 (2016).
- Ruda-Kucerova, J., Amchova, P., Siska, F., et al. NBQX attenuates relapse of nicotine seeking but not nicotine and methamphetamine self-administration in rats. World J. Biol. Psychiatry 22(10), 733-743 (2021).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 10/02/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM