



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## 9(S)-HODE

Item No. 38410

**CAS Registry No.:** 73543-67-6  
**Formal Name:** 9S-hydroxy-10E,12Z-octadecadienoic acid  
**MF:** C<sub>18</sub>H<sub>32</sub>O<sub>3</sub>  
**FW:** 296.5  
**Purity:** ≥98%  
**UV/Vis.:** λ<sub>max</sub>: 234 nm ε: 23,000  
**Supplied as:** A solution in ethanol  
**Storage:** -20°C  
**Stability:** ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

9(S)-HODE is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 9(S)-HODE in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 9(S)-HODE is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 9(S)-HODE in PBS (pH 7.2) is approximately 1 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

9(S)-HODE is produced by the lipoxygenation of linoleic acid in both plants and animals.<sup>1,2</sup> It has been detected in atherosclerotic plaques, as an esterified component of membrane phospholipids and in oxidized LDL particles.<sup>3</sup>

### References

1. Gardner, H.W. Soybean lipoxygenase-1 enzymically forms both (9S)- and (13S)-hydroperoxides from linoleic acid by a pH-dependent mechanism. *Biochim. Biophys. Acta* **1001(3)**, 274-281 (1989).
2. Kühn, H., Belkner, J., and Wiesner, R. Subcellular distribution of lipoxygenase products in rabbit reticulocyte membranes. *Eur. J. Biochem.* **191(1)**, 221-227 (1990).
3. Ku, G., Thomas, C.E., Akeson, A.L., *et al.* Induction of interleukin 1β expression from human peripheral blood monocyte-derived macrophages by 9-hydroxyoctadecadienoic acid. *J. Biol. Chem.* **267(20)**, 14183-14188 (1992).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/22/2021

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM