

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Malathion-d₆ Item No. 39262

CAS Registry No.: 1189877-72-2

Formal Name: 1,4-diethyl 2-[[di(methoxy-d₃)

phosphinothioyl]thio]butanedioate

MF: $C_{10}H_{13}D_6O_6PS_2$

FW: 336.4

Chemical Purity: ≥95% (Malathion)

Deuterium

 \geq 99% deuterated forms (d₁-d₆); \leq 1% d₀ Incorporation:

Supplied as: A solution in methanol

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Malathion-d₆ is intended for use as an internal standard for the quantification of malathion (Item No. 22998) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Malathion- d_{κ} is supplied as a solution in methanol. To change the solvent, simply evaporate the methanol under a gentle stream of nitrogen and immediately add the solvent of choice. Malathion-d₆ is slightly soluble in chloroform.

Description

Malathion is an organophosphate insecticide that inhibits acetylcholinesterase (AChE; $IC_{50}s = 370$ and 160 µM for free and immobilized AChE, respectively, from bovine erythrocytes). 1 It is ovicidal and adulticidal to laboratory colonies of body lice when used at concentrations of 0.001 and 0.05%, respectively.² It is toxic to rats (LD₅₀ = 750 mg/kg) and teratogenic to zebrafish when used at concentrations ranging from 2 to 3 mg/L.^{3,4} Formulations containing malathion have been used in the treatment of lice.

References

- 1. Krstić, D.Z., Ĉolović, M., Kralj, M.B., et al. Inhibition of AChE by malathion and some structurally similar compounds. J. Enzyme Inhib. Med. Chem. 23(4), 562-573 (2008).
- Cole, M.M. and Burden, G.S. Phosphorus compounds as ovicides and adulticides against body lice. J. Econ. Entomol. 49(6), 747-750 (1956).
- DuBois, K.P., Doull, J., Deroin, J., et al. Toxicity and mechanism of action of some new insecticidal thionophosphates. Arch. Ind. Hyg. Occup. Med. 8(4), 350-358 (1953).
- Cook, L.W., Paradise, C.J., and Lom, B. The pesticide malathion reduces survival and growth in developing zebrafish. Environ. Toxicol. Chem. 24(7), 1745-1750 (2005).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 07/12/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM