

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Carbofuran-d₃ Item No. 39558

CAS Registry No.: 1007459-98-4

Formal Name: N-(methyl-d₂)-carbamic acid, 2,3-dihydro-

2,2-dimethyl-7-benzofuranyl ester

Synonym: BAY 70143-d₃ MF: $C_{12}H_{12}D_3NO_3$

224.3 FW:


≥98% (Carbofuran) **Chemical Purity:**

Deuterium

 \geq 99% deuterated forms (d₁-d₃); \leq 1% d₀ Incorporation:

Supplied as: A solid -20°C Storage: Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Carbofuran-d₃ is intended for use as an internal standard for the quantification of carbofuran (Item No. 25635) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Carbofuran-d₃ is supplied as a solid. A stock solution may be made by dissolving the carbofuran-d₃ in the solvent of choice. Carbofuran-d₃ is slightly soluble in methanol (warm) and chloroform.

Description

Carbofuran is a carbamate insecticide that inhibits acetylcholinesterase (AChE) in insects and mammals (K_ss = 9.4, 30, and 210 μM for M. domestica brain, A. mellifera brain, and bovine erythrocyte AChE, respectively). It also inhibits AChE from rat and human erythrocytes with IC₅₀ values of 40 and 25 nM, respectively. It is toxic to the housefly M. domestica (LD₅₀ = 4.6 mg/kg) and fourth-instar larvae of the mosquito C. pipiens with an LC_{50} value of 0.054 ppm. Carbofuran is toxic to birds, rats (LD_{50} = 4-11 mg/kg), and other mammals, including dogs, sheep, and cattle, and is considered an environmental toxin.^{1,3} Formulations containing carbofuran have been used in agriculture for the control of insects, nematodes, mites, and ticks.

References

- 1. Yu, C., Metcalf, R.L., and Booth, G.M. Inhibition of acetylcholinesterase from mammals and insects by carbofuran and its related compounds and their toxicities toward these animals. J. Agric. Food Chem. **20(5)**, 923-926 (1972).
- 2. Rao, P.S., Roberts, G.H., Pope, C.N., et al. Comparative inhibition of rodent and human erythrocyte acetylcholinesterase by carbofuran and carbaryl. Pestic. Biochem. Physiol. 48(2), 79-84 (1994).
- 3. Gupta, R.C. Carbofuran toxicity. J. Toxicol. Environ. Health 43(4), 383-418 (1994).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/12/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM