



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



(R)-(-)-Metalaxyl-d₆

Item No. 39575

CAS Registry No.: 1398112-32-7
Formal Name: methyl N-(2,6-bis(methyl-d₃)phenyl)-N-(2-methoxyacetyl)-D-alaninate
Synonym: (R)-(-)-Metalaxyl-d₆
MF: C₁₅H₁₅D₆NO₄
FW: 285.4
Chemical Purity: ≥95% ((R)-(-)-metalaxyl)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₆); ≤1% d₀
Supplied as: A solution in acetonitrile
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(R)-(-)-Metalaxyl-d₆ is intended for use as an internal standard for the quantification of (R)-(-)-metalaxyl by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

(R)-(-)-Metalaxyl is a fungicide and an active enantiomer of metalaxyl (Item No. 25817).^{1,2} It is active against several isolates of the plant pathogenic fungus *P. nicotianae* (EC₅₀s = <0.01-2.9 µg/ml).¹ (R)-(-)-Metalaxyl (3 g/kg of seeds) completely protects sunflower plants (*H. annuus*) from infection by *P. helianthi*.² It is toxic to *D. magna* and *D. rerio* (LC₅₀s = 176.4 and 237.67 mg/L, respectively).³ Formulations containing (R)-(-)-metalaxyl have been used in the control of fungi in agriculture.

References

1. Hu, J., Hong, C., Moorman, G.W., *et al.* Mefenoxam sensitivity and fitness analysis of Phytophthora nicotianae isolates from nurseries in Virginia, USA. *Plant Pathol.* **57(4)**, 728-736 (2008).
2. Zadra, C., Marucchini, C., and Zizzerini, A. Behavior of metalaxyl and its pure R-enantiomer in sunflower plants (*Helianthus annuus*). *J. Agric. Food Chem.* **50(19)**, 5373-5377 (2002).
3. Yao, K., Zhu, L., Duan, Z., *et al.* Comparison of R-metalaxyl and rac-metalaxyl in acute, chronic, and sublethal effect on aquatic organisms: *Daphnia magna*, *Scenedesmus quadricanda*, and *Danio rerio*. *Environ. Toxicol.* **24(2)**, 148-156 (2009).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/19/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM