



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



Trifloxystrobin-d₆

Item No. 39591

Formal Name: methyl (E)-2-(methoxyimino)-2-(((((E)-1-(3-(trifluoromethyl)phenyl)ethylidene)amino)oxy)methyl-d₂)phenyl-3,4,5,6-d₄)acetate

MF: C₂₀H₁₃D₆F₃N₂O₄
FW: 414.4

Chemical Purity: ≥95% (Trifloxystrobin)


Deuterium

Incorporation: ≥99% deuterated forms (d₁-d₆); ≤1% d₀

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Trifloxystrobin-d₆ is intended for use as an internal standard for the quantification of trifloxystrobin (Item No. 25825) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Trifloxystrobin-d₆ is supplied as a solid. A stock solution may be made by dissolving the trifloxystrobin-d₆ in the solvent of choice, which should be purged with an inert gas. Trifloxystrobin-d₆ is slightly soluble in chloroform and methanol.

Description

Trifloxystrobin is a fungicide.^{1,2} It inhibits the growth of *R. solani* isolates from sugar beet crops (EC₅₀s = 0.14-823.54 µg/ml).¹ Trifloxystrobin (100-1,000 ppm) inhibits mycelial growth of *C. gloeosporioides* when used in combination with tebuconazole (Item No. 24052).² It is toxic to various aquatic species, including *R. arenarium*, *P. santafecinus*, *E. bicolor*, and *L. latrans* tadpoles (LC₅₀s = 0.22, 0.14, 0.1, and 0.26 mg/L, respectively).³ It also reduces mobility of tadpoles and decreases predation of trifloxystrobin-exposed, but not non-exposed, tadpoles by eels. Formulations containing trifloxystrobin have been used in the control of fungi in agriculture and aquatic areas.

References

- Arabiati, S. and Khan, M. Sensitivity of *Rhizoctonia solani* AG-2-2 from sugar beet to fungicides. *Plant Dis.* **100(12)**, 2427-2433 (2016).
- Dev, D. and Narendrappa, T. In vitro evaluation of fungicides against *Colletotrichum gloeosporioides* (Penz.) Penz and Sacc. causing anthracnose of pomegranate (*Punica granatum* L.). *J. Appl. Nat. Sci.* **8(4)**, 2268-2272 (2017).
- Junges, C.M., Peltzer, P.M., Lajmanovich, R.C., et al. Toxicity of the fungicide trifloxystrobin on tadpoles and its effect on fish-tadpole interaction. *Chemosphere* **87(11)**, 1348-1354 (2012).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM