



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## SAINT C-18

Item No. 39709

**CAS Registry No.:** 187587-97-9  
**Formal Name:** 1-methyl-4-[(10Z)-1-(9Z)-9-octadecen-1-yl-10-nonadecen-1-yl]-pyridinium, monochloride  
**Synonyms:** SAINT-2, SAINT-7, SAINT-18, Synthetic Amphiphile INTERaction C-18

**MF:** C<sub>43</sub>H<sub>78</sub>N • Cl


**FW:** 644.5

**Purity:** ≥98%

**Supplied as:** A solution in chloroform

**Storage:** -20°C

**Stability:** ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

SAINT C-18 is supplied as a solution in chloroform. To change the solvent, simply evaporate the chloroform under a gentle stream of nitrogen and immediately add the solvent of choice, which should be purged with an inert gas. SAINT C-18 is slightly soluble in acetonitrile.

### Description

SAINT C-18 is a cationic lipid that has been used in the formation of liposomes for the delivery of plasmid DNA, proteins, siRNA, and small molecules.<sup>1-4</sup> Complexes of SAINT C-18 with plasmid DNA encoding pigment epithelium-derived factor (PEDF) inhibit corneal neovascularization induced by complexes of SAINT C-18 with plasmid DNA encoding FGF2, also known as basic FGF, in rats.<sup>2</sup>

### References

1. van der Woude, I., Wagenaar, A., Meekel, A.A.P., *et al.* Novel pyridinium surfactants for efficient, nontoxic *in vitro* gene delivery. *Proc. Natl. Acad. Sci. USA* **94**(4), 1160-1165 (1997).
2. Kuo, C.-N., Yang, L.-C., Yang, C.-T., *et al.* Inhibition of corneal neovascularization with plasmid pigment epithelium-derived factor (p-PEDF) delivered by synthetic amphiphile INTERaction-18 (SAINT-18) vector in an experimental model of rat corneal angiogenesis. *Exp. Eye Res.* **89**(5), 678-685 (2009).
3. van der Gun, B.T.F., Monami, A., Laarmann, S., *et al.* Serum insensitive, intranuclear protein delivery by the multipurpose cationic lipid SAINT-2. *J. Control Release* **123**(3), 228-238 (2007).
4. Adrian, J.E., Morselt, H.W., Süß, R., *et al.* Targeted SAINT-O-Somes for improved intracellular delivery of siRNA and cytotoxic drugs into endothelial cells. *J. Control Release* **144**(3), 341-349 (2010).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/26/2024

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897

[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM