



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## Mepivacaine-d<sub>3</sub> Item No. 39812

**CAS Registry No.:** 1346597-90-7  
**Formal Name:** N-(2,6-dimethylphenyl)-1-(methyl-d<sub>3</sub>)  
piperidine-2-carboxamide  
**MF:** C<sub>15</sub>H<sub>19</sub>D<sub>3</sub>N<sub>2</sub>O  
**FW:** 249.4  
**Chemical Purity:** ≥98% (Mepivacaine)  
**Deuterium Incorporation:** ≥99% deuterated forms (d<sub>1</sub>-d<sub>3</sub>); ≤1% d<sub>0</sub>  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:** ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Mepivacaine-d<sub>3</sub> is intended for use as an internal standard for the quantification of mepivacaine by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Mepivacaine-d<sub>3</sub> is supplied as a solid. A stock solution may be made by dissolving the mepivacaine-d<sub>3</sub> in the solvent of choice, which should be purged with an inert gas. Mepivacaine-d<sub>3</sub> is slightly soluble in chloroform and methanol.

### Description

Mepivacaine is an inhibitor of voltage-gated sodium channels (Na<sub>v</sub>s) and a local anesthetic.<sup>1</sup> It inhibits depolarization-induced currents in ND7/23 cells expressing Na<sub>v</sub>1.8 channels when used at concentrations of 30 and 100 μM. Mepivacaine (2% w/v) reduces lameness in horses with naturally occurring forelimb lameness.<sup>2</sup> Formulations containing mepivacaine have been used as local and regional anesthetics.

### References

1. Leffler, A., Reckzeh, J., and Nau, C. Block of sensory neuronal Na<sup>+</sup> channels by the secretolytic ambroxol is associated with an interaction with local anesthetic binding sites. *Eur. J. Pharmacol.* **630(1-3)**, 19-28 (2010).
2. Boorman, S., DeGraves, F., Schumacher, J., et al. Comparison of 2% mepivacaine and a solution of 2% lidocaine/epinephrine administered for median and ulnar nerve blocks in horses with naturally occurring forelimb lameness. *Vet. Surg.* **51(2)**, 279-285 (2022).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/11/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM