



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## (+)-Fenfluramine (hydrochloride)

Item No. 39830

**CAS Registry No.:** 3239-45-0  
**Formal Name:** N-ethyl- $\alpha$ -methyl-3-(trifluoromethyl)-benzeneethanamine, monohydrochloride  
**Synonym:** Dexfenfluramine  
**MF:** C<sub>12</sub>H<sub>16</sub>F<sub>3</sub>N • HCl  
**FW:** 267.7  
**Purity:**  $\geq$ 95%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:**  $\geq$ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

(+)-Fenfluramine (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the (+)-fenfluramine (hydrochloride) in the solvent of choice, which should be purged with an inert gas. (+)-Fenfluramine (hydrochloride) is soluble in organic solvents such as DMSO. It is also soluble in water. We do not recommend storing the aqueous solution for more than one day.

### Description

(+)-Fenfluramine is an agonist of the serotonin (5-HT) receptor subtypes 5-HT<sub>2B</sub> and 5-HT<sub>2C</sub>.<sup>1</sup> It induces serotonin release from superfused midbrain synaptosome preparations isolated from wild-type mice, but not 5-HT<sub>2B</sub><sup>-/-</sup> mice, when used at concentrations of 0.5 or 1  $\mu$ M. (+)-Fenfluramine (3 or 10 mg/kg) reduces food intake in starved wild-type mice but not in starved 5-HT<sub>2B</sub><sup>-/-</sup> mice. It reduces body weight gain in wild-type mice when infused at a dose of 2.5 mg/kg per day for 35 days. Formulations containing (+)-fenfluramine were previously used in the treatment of obesity.

### Reference

1. Banas, S.M., Doly, S., Boutourlinsky, K., *et al.* Deconstructing antiobesity compound action: Requirement of serotonin 5-HT<sub>2B</sub> receptors for dexfenfluramine anorectic effects. *Neuropsychopharmacology* **36(2)**, 423-433 (2011).

**WARNING**  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

**SAFETY DATA**  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

**WARRANTY AND LIMITATION OF REMEDY**  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/18/2024

### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM