

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Bepotastine

Item No. 39932

CAS Registry No.: 125602-71-3

Formal Name: 4-[(S)-(4-chlorophenyl)-2-pyridinylmethoxy]-

1-piperidinebutanoic acid

MF: $C_{21}H_{25}CIN_2O_3$

FW: 388.9 **Purity:** ≥98% Supplied as: A solid Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Bepotastine is supplied as a solid. A stock solution may be made by dissolving the bepotastine in the solvent of choice, which should be purged with an inert gas. Bepotastine is soluble in organic solvents such as methanol and DMSO. It is also soluble in water. We do not recommend storing the aqueous solution for more than one day.

Description

Bepotastine is an antagonist of the histamine H_1 receptor that is selective over H_3 , α_1 -, α_2 -, and β-adrenergic, dopamine D_{2long} , serotonin 5-HT $_2$, muscarinic acetylcholine, and benzodiazepine receptors. It reduces dye leakage from the nasal passages of rats acutely sensitized to an antigen ($ED_{50} = 0.03 \text{ mg/kg}$) and inhibits histamine-induced bronchoconstriction in the anesthetized dog (ED $_{50}$ = 3.2 μ g/kg). 2,3 Bepotastine prevents conjunctival vascular hyperpermeability in a guinea pig model of conjunctivitis in a dose-dependent manner.⁴ Formulations containing bepotastine have been used in the treatment of itching associated with allergic conjunctivitis.

References

- 1. Kato, M., Nishida, A., Aga, Y., et al. Pharmacokinetic and pharmacodynamic evaluation of central effect of the novel antiallergic agent betotastine besilate. Arzneimittelforschung. 47(10), 1116-1124 (1997).
- 2. Matsubara, S., Ono, C., Yamazaki, N., et al. Inhibitory effects of bepotastine, a novel anti-allergic drug on histamine-induced bronchoconstriction in anesthetized dogs. Yakuri to Chiryo 25(4), 895-900 (1997).
- Murata, T., Matsumoto, Y., Suzuki, T., et al. Effect of betotastine besilate (TAU-284), a novel anti-allergic agent, on experimental allergic rhinitis. Arerugi 46(7), 576-584 (1997).
- 4. Kida, T., Fuji, A., Sakai, O., et al. Bepotastine besilate, a highly selective histamine H₁ receptor antagonist, suppresses vascular hyperpermeability and eosinophil recruitment in in vitro and in vivo experimental allergic conjunctivitis models. Exp. Eye Res. 91(1), 85-91 (2010).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 03/20/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM