



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




Soluble Fas Ligand/CD95 Ligand (human, recombinant; aa 134-281)
Item No. 40212

Overview and Properties

Synonyms: CD178, CD95L, FasL, FASLG, TNFL6, TNF Ligand Superfamily Member 6, TNFSF6
Source: Active recombinant human N-terminal His-tagged FasL expressed in yeast
Amino Acids: 134-281
Uniprot No.: P48023
Molecular Weight: 19.3 kDa
Storage: -80°C (as supplied)
Stability: ≥1 year
Purity: ≥90% estimated by SDS-PAGE
Supplied in: Lyophilized from sterile PBS, pH 7.4
Protein
Concentration: *batch specific* mg/ml
Activity: *batch specific* U/ml
Specific Activity: *batch specific* U/mg

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Images



Lane 1: MW Markers
Lane 2: Soluble Fas Ligand/CD95 Ligand

SDS-PAGE Analysis of Soluble Fas Ligand/CD95 Ligand. This protein has a calculated molecular weight of 19.3 kDa. It has an apparent molecular mass of approximately 26 kDa, by SDS-PAGE, under reducing conditions due to glycosylation.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.


SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/10/2023

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Immobilized Human Fas/CD95 hFc at 2 $\mu\text{g/ml}$ (100 $\mu\text{l/well}$) can bind Soluble Fas Ligand/CD95 Ligand (human, recombinant; aa 134-281), the EC_{50} of Soluble Fas Ligand/CD95 Ligand (human, recombinant; aa 134-281) is 40-200 ng/ml.

Description

Fas ligand (FasL), also known as CD95L, is a type II transmembrane protein and member of the TNF superfamily.¹ It is composed of a cytoplasmic domain, transmembrane domain, stalk region, TNF homology domain (THD), which is involved in homotrimerization, and Fas-binding domain. FasL is expressed in activated T cells, as well as the placenta, testis, lung, prostate, esophagus, uterus and spleen.² Binding of FasL to its receptor Fas leads to formation of a death-inducing signaling complex (DISC) composed of CD95 oligomers, the Fas-associated death domain protein (FADD), and procaspase-8, procaspase-10, and cellular FLICE-like inhibitory protein (c-FLIP), which induces apoptosis.³ Soluble FasL (sFasL) is produced by proteolytic cleavage of FasL by various matrix metalloproteinases (MMPs) and either inhibits FasL-induced apoptosis or induces apoptosis in a context-dependent manner.^{1,4} sFasL induces migration of isolated human neutrophils and apoptosis in primary human distal lung epithelial cells.^{5,6} It decreases ankylosis in a rat model of collagen-induced arthritis.⁷ Serum levels of sFasL are increased in patients with insulin-dependent diabetes mellitus, systemic lupus erythematosus (SLE), rheumatoid arthritis, gastric cancer, or leukemia. Cayman's Soluble Fas Ligand/CD95 Ligand (human, recombinant; aa 134-281) protein can be used for binding assays. This protein consists of 168 amino acids and has a calculated molecular weight of 19.3 kDa. By SDS-PAGE, under reducing conditions, the apparent molecular mass of the protein is 26 kDa due to glycosylation.

References

1. Levoine, N., Jean, M., and Legembre, P. CD95 structure, aggregation and cell signaling. *Front. Cell Dev. Biol.* **8**, 314 (2020).
2. Xerri, L., Devilard, E., Hassoun, J., *et al.* Fas ligand is not only expressed in immune privileged human organs but is also coexpressed with Fas in various epithelial tissues. *Mol. Pathol.* **50(2)**, 87-91 (1997).
3. Peter, M.E., Hadji, A., Murmann, A.E., *et al.* The role of CD95 and CD95 ligand in cancer. *Cell Death Differ.* **22(4)**, 549-559 (2015).
4. Suda, T., Hashimoto, H., Tanaka, M., *et al.* Membrane Fas ligand kills human peripheral blood T lymphocytes, and soluble Fas ligand blocks the killing. *J. Exp. Med.* **186(12)**, 2045-2050 (1997).
5. Ottonello, L., Tortolina, G., Amelotti, M., *et al.* Soluble Fas ligand is chemotactic for human neutrophilic polymorphonuclear leukocytes. *J. Immunol.* **162(6)**, 3601-3606 (1999).
6. Matute-Bello, G., Liles, W.C., Steinberg, K.P., *et al.* Soluble Fas ligand induces epithelial cell apoptosis in humans with acute lung injury (ARDS). *J. Immunol.* **163(4)**, 2217-2225 (1999).
7. Li, N.-L., Nie, H., Yu, Q.-W., *et al.* Role of soluble Fas ligand in autoimmune diseases. *World J. Gastroenterol.* **10(21)**, 3151-3156 (2004).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM