



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



Rucaparib (camsylate)

Item No. 40218

CAS Registry No.: 1859053-21-6

Formal Name: (1S,4R)-7,7-dimethyl-2-oxo-bicyclo[2.2.1]heptane-1-methanesulfonic acid, compd. with 8-fluoro-1,3,4,5-tetrahydro-2-[4-[(methylamino)methyl]phenyl]-6H-pyrrolo[4,3,2-ef][2]benzazepin-6-one (1:1)

Synonyms: AG-014699, PF-01367338

MF: C₁₉H₁₈FN₃O • C₁₀H₁₆O₄S


FW: 555.7

Purity: ≥98%

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Rucaparib (camsylate) is supplied as a solid. A stock solution may be made by dissolving the rucaparib (camsylate) in the solvent of choice, which should be purged with an inert gas. Rucaparib (camsylate) is soluble in methanol. Rucaparib (camsylate) is slightly soluble in acetonitrile and water. We do not recommend storing the aqueous solution for more than one day.

Description

Rucaparib is a poly(ADP-ribose) polymerase 1 (PARP1) inhibitor.¹ It reduces PARP1 activity in D283 Med medulloblastoma cells by 91.7% when used at a concentration of 1 μM. Rucaparib selectively inhibits the proliferation of MDA-MB-436 breast cancer cells expressing mutant BRCA1 (IC_{50} s = 1.3 μM) over MCF-7 breast cancer cells expressing wild-type BRCA1 and BRCA2 (IC_{50} s = 20.2 μM).² It potentiates the growth inhibition induced by the DNA alkylating agent prodrug temozolomide (Item No. 14163) in D283 Med and D384 Med medulloblastoma cells when used at a concentration of 0.4 μM.¹ In vivo, rucaparib (1 mg/kg per day) potentiates temozolomide-induced delays in tumor growth in D283 Med and D384 Med mouse xenograft models. Formulations containing rucaparib have been used in the treatment of BRCA mutation-associated, drug-resistant ovarian, fallopian tube, or primary peritoneal cancers.

References

- Daniel, R.A., Rozanska, A.L., Mulligan, E.A., et al. Central nervous system penetration and enhancement of temozolomide activity in childhood medulloblastoma models by poly(ADP-ribose) polymerase inhibitor AG-014699. *Br. J. Cancer* **103**(10), 1588-1596 (2010).
- Drew, Y., Mulligan, E.A., Vong, W.T., et al. Therapeutic potential of poly(ADP-ribose) polymerase inhibitor AG014699 in human cancers with mutated or methylated BRCA1 or BRCA2. *JNCI* **103**(4), 334-346 (2011).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM