



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic




PRODUCT INFORMATION



Ochratoxin C- $^{13}\text{C}_{20}$

Item No. 40239

Formal Name:	ethyl ((R)-5-chloro-8-hydroxy-3-(methyl- ^{13}C)-1-oxoisochromane-7-carbonyl- $^{13}\text{C}_{10}$)-L-phenylalaninate- $^{13}\text{C}_9$
Synonyms:	Ochratoxin A- $^{13}\text{C}_{20}$ ethyl ester, OTC- $^{13}\text{C}_{20}$, OTA- $^{13}\text{C}_{20}$ ethyl ester
MF:	$\text{C}_2[\text{^{13}C}]_{20}\text{H}_{22}\text{ClNO}_6$
FW:	451.7
Purity:	$\geq 98\%$
Supplied as:	A solution in methanol
Storage:	-20°C
Stability:	≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Description

OTC- $^{13}\text{C}_{20}$ is intended for use as an internal standard for the quantification of ochratoxin C (Item No. 20183) by GC- or LC-MS. OTC is a mycotoxin that has been found in *Aspergillus* and is an active metabolite of OTA (Item No. 11439).^{1,2} It is formed from OTA by gut microbiota, however, OTA can also be formed from OTC *in vivo*.^{2,3} OTC inhibits the proliferation and differentiation of THP-1 monocytes (IC_{50} s = 126 and 91 ng/ml, respectively) and induces apoptosis in the same cells when used at concentrations of 100 or 1,000 ng/ml.¹ It induces mortality in zebrafish embryos (LC_{50} = 0.32 nM).⁴ OTC has been found in red wine.⁵

References

1. Müller, G., Rosner, H., Rohrmann, B., et al. Effects of the mycotoxin ochratoxin A and some of its metabolites on the human cell line THP-1. *Toxicology* **184**(1), 69-82 (2003).
2. Galtier, P. and Alvinerie, M. *In vitro* transformation of ochratoxin A by animal microbial floras. *Ann. Rech. Vet.* **7**(1), 91-98 (1976).
3. Fuchs, R., Hult, K., Peraica, M., et al. Conversion of ochratoxin C into ochratoxin A *in vivo*. *Appl. Environ. Microbiol.* **48**(1), 41-42 (1984).
4. Csenki, Z., Garai, E., Faisal, Z., et al. The individual and combined effects of ochratoxin A with citrinin and their metabolites (ochratoxin B, ochratoxin C, and dihydrocitrinone) on 2D/3D cell cultures, and zebrafish embryo models. *Food Chem. Toxicol.* **158**, 112674 (2021).
5. Remiro, R., Ibáñez-Vea, M., González-Peñas, E., et al. Validation of a liquid chromatography method for the simultaneous quantification of ochratoxin A and its analogues in red wines. *J. Chromatogr. A* **1217**(52), 8249-8256 (2010).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM