



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## DCLK1-IN-1 Item No. 40448

**CAS Registry No.:** 2222635-15-4  
**Formal Name:** 5,11-dihydro-2-[[2-methoxy-4-(4-methyl-1-piperazinyl)phenyl]amino]-11-methyl-5-(2,2,2-trifluoroethyl)-6H-pyrimido[4,5-b][1,4]benzodiazepin-6-one  
**Synonyms:** DCLK1 Inhibitor 1,  
Doublecortin-like Kinase 1-IN-1,  
Doublecortin-like Kinase 1 Inhibitor 1

**MF:** C<sub>26</sub>H<sub>28</sub>F<sub>3</sub>N<sub>7</sub>O<sub>2</sub>  
**FW:** 527.5  
**Purity:** ≥95%  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:** ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

DCLK1-IN-1 is supplied as a solid. A stock solution may be made by dissolving the DCLK1-IN-1 in the solvent of choice, which should be purged with an inert gas. DCLK1-IN-1 is soluble in acetonitrile and DMSO.

### Description

DCLK1-IN-1 is an inhibitor of doublecortin-like kinase 1 (DCLK1; IC<sub>50</sub> = 17 nM).<sup>1</sup> It is selective for DCLK1 over leucine-rich repeat kinase 2 (LRRK2), bromodomain-containing protein 4 (BRD4), and ERK5 (IC<sub>50</sub>s = 7,000, >10,000, and 3,350 nM, respectively). DCLK1-IN-1 (1 μM) reduces the number of cells in primary patient-derived pancreatic cancer organoids. *In vivo*, DCLK1-IN-1 (10 mg/kg every other day) reduces plaque area and collagen levels in the aortic arches and serum levels of Tnf-α and Il-6 in an ApoE<sup>-/-</sup> mouse model of atherosclerosis induced by a high-fat diet.<sup>2</sup>

### References

1. Ferguson, F.M., Liu, Y., Harshbarger, W., *et al.* Synthesis and structure-activity relationships of DCLK1 kinase inhibitors based on a 5,11-Dihydro-6H-benzo[e]pyrimido[5,4-b][1,4]diazepin-6-one scaffold. *J. Med. Chem.* **63**(14), 7817-7826 (2020).
2. Huang, Z., Shen, S., Han, X., *et al.* Macrophage DCLK1 promotes atherosclerosis via binding to IKKβ and inducing inflammatory responses. *EMBO Mol. Med.* **15**(5), e17198 (2023).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/29/2024

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM