SZABO SCANDIC

Part of Europa Biosite

Produktinformation

- Forschungsprodukte & Biochemikalien
- Zellkultur & Verbrauchsmaterial
- Diagnostik & molekulare Diagnostik
- Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!

Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic

PRODUCT INFORMATION

oxLig-1 (technical grade)

Item No. 40740

CAS Registry No.:	352523-18-3
Formal Name:	3β-[(8-carboxy-1-oxooctyl)oxy]-cholest-5-en-7-one
Synonym:	7-Ketocholesteryl-9-carboxynonanoate
MF:	$C_{36}H_{58}O_5$
FW:	570.8
Purity:	≥80%
Supplied as:	A solid
Storage:	-20°C
Stability:	≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

oxLig-1 (technical grade) is supplied as a solid. A stock solution may be made by dissolving the oxLig-1 (technical grade) in the solvent of choice, which should be purged with an inert gas. oxLig-1 (technical grade) is slightly soluble (0.1-1 mg/ml) in DMSO.

Description

oxLig-1 is an ω-carboxylated 7-ketocholesteryl ester and a lipid component of oxidized LDL (oxLDL).[1] It binds to the ligand-binding domain of peroxisome proliferator-activated receptor γ (PPARγ), an effect that can be reversed by the PPARγ agonist troglitazone (Item No. 71750), and also binds to CD36, also known as scavenger receptor B2, when used at a concentration of 50 µg/ml.[2,3] oxLig-1 (20 µg/ml) increases cholesterol efflux and protein levels of liver X receptor α (LXRα) and ATP-binding cassette transporter 1 subfamily A (ABCA1) in THP-1 macrophages. Liposomes containing oxLig-1 and complexed with apolipoprotein H, also known as β_2-glycoprotein I (β_2-GPI), bind to an anti-apolipoprotein H antibody, which mediates uptake by J774A.1 macrophages.[1] oxLig-1 decreases intracellular lipid accumulation induced by oleic acid (Item Nos. 90260 | 24659) in HepG2 cells, as well as hepatic lipid accumulation in mice fed a high-fat diet.[3] Serum levels of IgG autoantibodies targeting a complex of oxLig-1 and apolipoprotein H are increased in patients with antiphospholipid syndrome with a history of arterial thrombosis.[4]

References

1. Kobayashi, K., Matsuura, E., Liu, Q., *et al.* A specific ligand for β2-glycoprotein I mediates autoantibody-dependent uptake of oxidized low density lipoprotein by macrophages. *J. Lipid Res.* **42(5)**, 697-709 (2001).
2. Chi, Y., Wang, L., Liu, Y., *et al.* 7-ketocholesteryl-9-carboxynonanoate enhances ATP binding cassette transporter A1 expression mediated by PPARγ in THP-1 macrophages. *Atherosclerosis* **234(2)**, 461-468 (2014).
3. Fu, C., Xiang, M.-L., Chen, S., *et al.* Molecular drug simulation and experimental validation of the CD36 receptor competitively binding to long-chain fatty acids by 7-ketocholesteryl-9-carboxynonanoate. *ACS Omega* **8(31)**, 28277-28289 (2023).
4. Lopez, D., Kobayashi, K., Merrill, J.T., *et al.* IgG autoantibodies against β2-glycoprotein I complexed with a lipid ligand derived from oxidized low-density lipoprotein are associated with arterial thrombosis in antiphospholipid syndrome. *Clin. Dev. Immunol.* **10(2-4)**, 203-211 (2003).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.**CAYMANCHEM**.COM