



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## Cdk2 Isoform 1 (human, recombinant)


Item No. 40793

### Overview and Properties

**Synonyms:** Cell Division Protein Kinase 2, Cyclin-dependent Kinase 2, p33 Protein Kinase  
**Source:** Recombinant human C-terminal His-tagged Cdk2 isoform 1 expressed in insect cells  
**Amino Acids:** 1-298 (full length)  
**Peptide Sequence:**  
**Uniprot No.:** P24941  
**Molecular Weight:** 35 kDa  
**Storage:** -80°C (as supplied)  
**Stability:** ≥1 year  
**Purity:** ≥85% estimated by SDS-PAGE, ≥95% determined by SEC-HPLC  
**Supplied in:** Lyophilized from sterile 50 mM Tris, pH 8.0, with 100 mM sodium chloride and 10% glycerol  
**Endotoxin Testing:** <1.0 EU/μg, determined by the LAL endotoxin assay  
**Bioactivity:** See figures for details


Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Images



Lane 1: MW Markers  
Lane 2: Cdk2 Isoform 1

**SDS-PAGE Analysis of Cdk2 Isoform 1.** This protein has a calculated molecular weight of 35 kDa.



**SEC-HPLC verification of Cdk2 isoform 1.** This protein has a purity of ≥95%.

**WARNING**  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

**SAFETY DATA**  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

**WARRANTY AND LIMITATION OF REMEDY**  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/16/2024

**CAYMAN CHEMICAL**  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## Description

---

Cyclin-dependent kinase 2 (Cdk2) is a serine/threonine kinase and member of the CDK family of kinases involved in cell cycle regulation.<sup>1,2</sup> It is composed of an N-terminal domain, an activation segment, also known as the T loop, and a catalytic C-terminal domain.<sup>2</sup> Alternative splicing of *CDK2* mRNA produces a full-length isoform or a shorter isoform that lacks exon 5, which encodes for part of the Cdk2 activation segment and contains residues that participate in interactions with cyclins.<sup>3</sup> It is ubiquitously expressed and localizes to the cytoplasm or nucleus depending on the cell cycle stage.<sup>5,6</sup> Cdk2 associates with cyclin E to mediate the G<sub>1</sub>/S transition and with cyclin A during S phase progression.<sup>1,2</sup> Knockout of *Cdk2* induces gametogenesis defects in mice but does not affect mitotic cell division.<sup>3</sup> Protein levels of Cdk2 are increased in patients with oral squamous cell carcinoma, and overexpression of Cdk2/cyclin A or Cdk2/cyclin E is associated with poor prognosis in patients with oral cancer.<sup>7</sup> Cayman's Cdk2 Isoform 1 (human, recombinant) protein consists of 308 amino acids and has a calculated molecular weight of 35 kDa.

## References

---

1. Funk, J.O. Cell cycle checkpoint genes and cancer. *Encyclopedia of life sciences*. John Wiley & Sons (2005).
2. Li, Y., Zhang, J., Gao, W., *et al.* Insights on structural characteristics and ligand binding mechanisms of CDK2. *Int. J. Mol. Sci.* **16(5)**, 9314-9340 (2015).
3. Ortega, S., Prieto, I., Odajima, J., *et al.* Cyclin-dependent kinase 2 is essential for meiosis but not for mitotic cell division in mice. *Nat. Genet.* **35(1)**, 25-31 (2003).
4. Ji, X., Humenik, J., Yang, D., *et al.* PolyC-binding proteins enhance expression of the CDK2 cell cycle regulatory protein *via* alternative splicing. *Nucleic Acids Res.* **46(4)**, 2030-2044 (2018).
5. Sherr, C.J. and Roberts, J.M. Living with or without cyclins and cyclin-dependent kinase. *Genes Dev.* **18(22)**, 2699-2711 (2004).
6. Bresnahan, W.A., Thompson, E.A., and Albrecht, T. Human cytomegalovirus infection results in altered Cdk2 subcellular localization. *J. Gen. Virol.* **78(Pt 8)**, 1993-1997 (1997).
7. Mihara, M., Shintani, S., Nakahara, Y., *et al.* Overexpression of CDK2 is a prognostic indicator of oral cancer progression. *Jpn. J. Cancer Res.* **92(3)**, 352-360 (2001).

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM