



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



LMT-28

Item No. 40837

CAS Registry No.: 1239600-18-0
Formal Name: (4S)-3-[(2S,3S)-3-hydroxy-2-methyl-4-methylene-1-oxononyl]-4-(1-methylethyl)-2-oxazolidinone
MF: C₁₇H₂₉NO₄
FW: 311.4
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

LMT-28 is supplied as a solid. A stock solution may be made by dissolving the LMT-28 in the solvent of choice, which should be purged with an inert gas. LMT-28 is soluble (≥10 mg/ml) in ethanol and DMSO.

Description

LMT-28 is an inhibitor of the protein-protein interaction between glycoprotein 130 (gp130) and the IL-6-bound form of IL-6 receptor subunit α (IL-6R α).¹ It binds to gp130 ($K_d = 7.4 \mu\text{M}$) and selectively inhibits IL-6-induced STAT3 phosphorylation over leukemia inhibitor factor- or phorbol 12-myristate 13-acetate-induced STAT3 phosphorylation in HepG2 cells at 10, 30, or 100 μM . LMT-28 (10 to 1,000 nM) reduces IL-6 induced increases in the viability of GM-CSF-starved TF-1 leukemia cells. It inhibits the increases in mRNA encoding Rankl, as well as inhibits increases in the phosphorylation levels of gp130, Stat3, and JAK2, induced by LPS and advanced glycation end products (AGEs) in mouse MC3T3-E1 osteoblasts when used at a concentration of 100 nM.² LMT-28 (0.25 mg/kg per day) decreases joint and paw swelling, as well as serum levels of cartilage oligomeric matrix protein (COMP), serum amyloid protein (SAP), and anti-collagen II antibodies, in a mouse model of collagen-induced arthritis.¹ Oral administration of LMT-28 (0.23 mg/kg per day) reduces bone resorption height and volume in a rat model of type 2 diabetes peri-implantitis induced by LPS, *P. gingivalis*, and a high-fat and high-sugar diet.²

References

- Hong, S.-S., Choi, J.H., Lee, S.Y., *et al.* A novel small-molecule inhibitor targeting the IL-6 receptor β subunit, glycoprotein 130. *J. Immunol.* **195**(1), 237-245 (2015).
- Liu, Q.-Q., Wu, W.-W., Yang, J., *et al.* A GP130-targeting small molecule, LMT-28, reduces LPS-induced bone resorption around implants in diabetic models by inhibiting IL-6/GP130/JAK2/STAT3 signaling. *Mediators Inflamm.* 933049 (2023).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/10/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM