

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Product Information

Prostaglandin E₂-MOX AChE Tracer

Catalog No. 414730

Sold under license SPI-BIO



IMPORTANT

Water used to prepare all EIA reagents and buffers must be deionized and free of trace organic contaminants ("UltraPure"). Use activated carbon filter cartridges or other organic scavengers. Glass distilled water (even if double distilled), HPLC-grade water, and sterile water (for injections) are not adequate for EIA. UltraPure water may also be purchased (Catalog No. 400000).

Laboratory Procedures

This vial contains lyophilized Prostaglandin E2-MOX Acetylcholinesterase (PGE2-MOX AChE) EIA Tracer (a covalent conjugate of PGE2-MOX and electric eel AChE (EC 3.1.1.7)). For long term storage, we suggest that the tracer be stored as supplied at -20°C; it will be stable for at least one year. Reconstitute 100 determination vials with 6 ml EIA Buffer or 500 determination vials with 30 ml EIA Buffer (see buffer preparation instructions below). Store the reconstituted tracer at 4°C and use within one week. For your convenience, we have supplied a 20% surplus of tracer.

Sample and Standard Preparation

This assay measures the methoximated derivative of PGE₂ (PGE₂-MOX). All standards and samples must be methoximated prior to performing the assay. See the methoximation protocol on the following page for details.

Buffer Preparation

1. Phosphate Buffer

Prepare a 1.0 M phosphate buffer solution by combining 133 g K₂HPO₄ and 32.15 g KH₂PO₄ and diluting to a total volume of 1.0 liter with UltraPure water. The pH of this solution will be 7.4.

Combine 100 ml of the phosphate buffer prepared above with 100 mg NaN₃, 23.4 g NaCl, 370 mg tetrasodium EDTA, and 1 g bovine serum albumin (Sigma A7030 or equivalent). Stir at room temperature until completely dissolved and dilute to a total volume of 1.0 liter with UltraPure water. Add 0.1 ml Tween 20 and mix thoroughly. This EIA Buffer may also be purchased from Cayman as a 10X concentrate buffer (Catalog No. 400060).

Combine 10 ml of the phosphate buffer prepared above with 0.5 ml Tween 20. Bring to a final volume of 1.0 liter with UltraPure water. This Wash Buffer may also be purchased from Cayman as a 400X concentrate buffer (Catalog No. 400062).

Methoximation Protocol

Prepare the methyloximating reagent by adding 0.82 g sodium acetate and 0.1 g methoxylamine HCl to 10 ml of a 10:90 ethanol/water solution (both sodium acetate and methoxylamine are available from Cayman Chemical, see front for catalog numbers). This reagent is not stable so it should all be used at one time.

Heat a 1:1 solution of PGE2 standard or sample and the methoximating reagent at 60°C for 30 minutes. Centrifuge the samples (standards do not need to be centrifuged) and dilute the standard or sample supernatant at least 1:5 with EIA Buffer before addition to the assay well.

WARNING: This product is for laboratory research only: not for administration to humans. Not for human or veterinary DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, or inhale. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Before use, the user must review the complete Material Safety Data Sheet, which has been sent under separate cover to the MSDS supervisor at your institution.

Cayman Chemical Company makes **no warranty or guarantee** of any kind, whether written or oral, expressed or implied, including without limitation, any warranty of fitness for a particular ose, suitability and merchantability, which extends beyond the description of the chemicals hereof. Cayman warrants only to the original customer that the material will meet our specifications

Target and the time of delivery.

Cayman will carry out its delivery obligations with due care and skill. Thus, in no event will Cayman have **any obligation or liability**, whether in tort (including negligence) or in contract, for

direct, indirect, incidental or consequential damages, even if Cayman is informed about their possible existence.

This limitation of liability does not apply in the case of intentional acts or negligence of Cayman, its directors or its employees.

Buyer's exclusive remedy and Cayman's sole liability hereunder shall be limited to a refund of the purchase price, or at Cayman's option, the replacement, at no cost to Buyer, of all material that

does not meet our specifications.
Said refund or replacement is conditioned on Buyer giving written notice to Cayman within thirty (30) days after arrival of the material at its destination. Failure of Buyer to give said notice within thirty (30) days shall constitute a waiver by Buyer of all claims hereunder with respect to said material.

For further details, please refer to our Warranty and Limitation of Remedy located on our website and in our catalog.

Copyright Cayman Chemical Company, 06/16/2009

Cayman Chemical

Mailing address

1180 E. Ellsworth Road Ann Arbor, MI 48108 USA

(800) 364-9897 (734) 971-3335

(734) 971-3640

custserv@caymanchem.com

www.caymanchem.com

Product Information

Suggested Assay Protocol

This tracer has been tested and formulated to work exclusively with ACETM reagents. This tracer may not perform as described if used with other assay reagents or protocols. *NOTE: this is an abbreviated protocol. If you are not familiar with this assay, please contact us for a complete protocol.*

- 1. Add 100 μ l of EIA Buffer to NSB wells and 50 μ l to B₀ wells.
- 2. Add 50 µl of Standard/sample to the appropriate wells.
- 3. Add 50 µl Tracer to all wells except Blk and TA.
- 4. Add 50 μl Antiserum (Catalog No. 414732) to all wells except Blk, TA, and NSB.
- 5. Incubate overnight at 4°C.
- 6. Wash the plate five times with Wash Buffer.
- 7. Add 200 µl Ellman's Reagent to each well.
- 8. Add 5 µl Tracer to the TA well.
- 9. Develop for approximately 90-120 minutes ($B_0 = 0.3-0.8 \text{ AU}$).
- 10. Read absorbance at a wavelength between 405 and 420 nm.

Blk-Blank; **NSB**-Non-specific Binding; **B**₀-Maximum Binding; **TA**-Total Activity;

S1-S8-Standards; 1-8-Samples

3

Related Products

Prostaglandin E $_2$ - Cat. No. 14010 • UltraPure Water - Cat. No. 400000 • Mouse Monoclonal Anti-Rabbit IgG - Cat. No. 400003 • Precoated (Mouse Anti-Rabbit IgG) EIA 96-Well Strip Plates - Cat. No. 400005 • Precoated (Mouse Anti-Rabbit IgG) EIA 96-Well Plates - Cat. No. 400007 • 96-Well Cover Sheets - Cat. No. 400012 • SPE Cartridges (C-18) - Cat. No. 400020 • Tween 20 - Cat. No. 400035 • Methoxylamine HCl - Cat. No. 400036 • Sodium Acetate - Cat. No. 400037 • Ellman's Reagent - Cat. No. 400050 • EIA Buffer Concentrate (10X) - Cat. No. 400060 • Wash Buffer Concentrate (400X) - Cat. No. 400062 • Prostaglandin E $_2$ -MOX EIA Antiserum - Cat. No. 414732

Cayman Chemical

Mailing address

1180 E. Ellsworth Road Ann Arbor, MI 48108 USA

Phone

(800) 364-9897 (734) 971-3335

Fax

(734) 971-3640

-Mail

custserv@caymanchem.com

Web

www.caymanchem.com