



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Tocainide

Item No. 41592

CAS Registry No.: 41708-72-9
Formal Name: 2-amino-N-(2,6-dimethylphenyl)-propanamide
Synonyms: (R,S)-Tocainide, W-36095
MF: C₁₁H₁₆N₂O
FW: 192.3
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Tocainide is supplied as a solid. A stock solution may be made by dissolving the tocainide in the solvent of choice, which should be purged with an inert gas. Tocainide is slightly soluble (0.1-1 mg/ml) in ethanol and soluble (≥10 mg/ml) in DMSO.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of tocainide can be prepared by directly dissolving the solid in aqueous buffers. The solubility of Tocainide is sparingly soluble (1-10 mg/ml) in PBS (pH 7.2). We do not recommend storing the aqueous solution for more than one day.

Description

Tocainide is an antiarrhythmic agent and a derivative of the local anesthetic lidocaine.¹ It inhibits voltage-gated sodium channel 1.4 (Na_v1.4) in HEK293 cells expressing the human channel (IC₅₀s = 699 and 182 μM at 0.1 and 10 Hz, respectively).² Tocainide reduces the action potential duration at 50% repolarization in isolated dog Purkinje fibers in a concentration-dependent manner.¹ *In vivo*, tocainide reduces ventricular heart rate and the percent of ectopic beats in a dog model of ventricular tachycardia induced by the cardiac glycoside ouabain (Item No. 14319).¹ Tocainide (50 mg/kg) decreases mechanical hyperalgesia in the nerve-injured paw in a rat model of neuropathic pain induced by chronic constriction injury of the sciatic nerve.² Formulations containing tocainide have previously been used in the treatment of ventricular arrhythmias.

References

1. Moore, E.N., Spear, J.F., Horowitz, L.N., *et al.* Electrophysiologic properties of a new antiarrhythmic drug — tocainide. *Am. J. Cardiol.* **41**(4), 703-709 (1978).
2. Ghelardini, C., J.-F., D., Muraglia, M., *et al.* Effects of a new potent analog of tocainide on hNav1.7 sodium channels and *in vivo* neuropathic pain models. *Neuroscience* **169**(2), 863-873 (2010).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/30/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM