



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRODUCT INFORMATION



## GDF8 Signaling Domain (human, recombinant)


Item No. 42188

### Overview and Properties

**Synonyms:** Growth Differentiation Factor 8, Myostatin  
**Source:** Recombinant human N-terminal human IgG1 FC-tagged GDF8 signaling domain expressed in HEK293 cells  
**Amino Acids:** 267-375  
**Uniprot No.:** O08689  
**Molecular Weight:** 40.8 kDa  
**Storage:** -80°C (as supplied)  
**Stability:** ≥1 year  
**Purity:** ≥92% estimated by SDS-PAGE  
**Supplied in:** Lyophilized from sterile PBS, pH 7.4  
**Endotoxin Testing:** <1.0 EU/μg, determined by the LAL endotoxin assay  
**Protein Concentration:** *batch specific* mg/ml

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Image



Lane 1: MW Markers  
Lane 2: GDF8 Signaling Domain

**SDS-PAGE Analysis of GDF8 Signaling Domain.** This protein has a calculated molecular weight of 40.8 kDa. It has an apparent molecular weight of approximately 45 kDa by SDS-PAGE under reducing conditions due to glycosylation.

WARNING  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/16/2024

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM

# PRODUCT INFORMATION



## Description

---

Growth differentiation factor 8 (GDF8) is a myokine and member of the TGF- $\beta$  superfamily.<sup>1,2</sup> It is produced in skeletal and cardiac muscle cells as a proprotein, which shares sequence identity between the human, mouse, and rat.<sup>1,5</sup> The proprotein is cleaved in the endoplasmic reticulum into a protein containing a prodomain and C-terminal signaling domain.<sup>1,3,5</sup> This protein is secreted and forms homodimers that either circulate or remain near the extracellular membrane in a latent state.<sup>6</sup> The prodomain is cleaved by bone morphogenetic protein 1 (BMP1) or tolloid family metalloproteinases, releasing the signaling domain to activate TGF- $\beta$  receptor type 2 (TGFB2) and negatively regulate muscle growth and differentiation.<sup>1,2,6</sup> Knockout of *Myst*, the gene expressing *Gdf8*, increases muscle mass, muscle fiber length, bone mineral content (BMC), and bone size in *Myst*<sup>-/-</sup> mice compared to wild-type mice.<sup>4</sup> The levels of full-length GDF8 and GDF8 prodomain are increased in left ventricle tissue samples from patients with dilated- or ischemic cardiomyopathy.<sup>5</sup> Cayman's GDF8 Signaling Domain (human, recombinant) protein is a disulfide-linked homodimer. The reduced monomer, composed of GDF8 (amino acids 267-375) fused to human IgG1 Fc at its N-terminus, consists of 369 amino acids and has a calculated molecular weight of 40.8 kDa. This protein consists of 108 amino acids and has a predicted N-terminus of Gly20 after signal peptide cleavage. As a result of glycosylation, the monomer migrates to approximately 45 kDa by SDS-PAGE under reducing conditions.

## References

---

1. Esposito, P., Picciotto, D., Battaglia, Y., *et al.* Myostatin: Basic biology to clinical application. *Adv. Clin. Chem.* **106**, 181-234 (2022).
2. McPherron, A.C., Lawler, A.M., and Lee, S.J. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* **387(6628)**, 83-90 (1997).
3. Sharma, M., Kambadur, R., Matthews, K.G., *et al.* Myostatin, a transforming growth factor-beta superfamily member, is expressed in heart muscle and is upregulated in cardiomyocytes after infarct. *J. Cell Physiol.* **180(1)**, 1-9 (1999).
4. Elkasrawy, M.N., and Hamrick, M.W. Myostatin (GDF-8) as a key factor linking muscle mass and skeletal form. *J. Musculoskelet. Neuronal. Interact.* **10(1)**, 56-63 (2010).
5. George, I., Bish, L.T., Kamalakkannan, G., *et al.* Myostatin activation in patients with advanced heart failure and after mechanical unloading. *Eur. J. Heart Fail.* **12(5)**, 444-453 (2010).
6. Wolfman, N.M., McPherron, A.C., Pappano, W.N., *et al.* Activation of latent myostatin by the BMP-1/tolloid family of metalloproteinases. *Proc. Natl. Acad. Sci. USA* **100(26)**, 15842-15846 (2003).

CAYMAN CHEMICAL  
1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA  
PHONE: [800] 364-9897  
[734] 971-3335  
FAX: [734] 971-3640  
CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM