



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Information



Maturation-Inducing Steroid (salmonid) EIA Antiserum

Item No. 498502

IMPORTANT

Water used to prepare all EIA reagents and buffers must be deionized and free of trace organic contaminants ('UltraPure'). Use activated carbon filter cartridges or other organic scavengers. Glass distilled water (even if double distilled), HPLC-grade water, and sterile water (for injections) are not adequate for EIA. UltraPure water may also be purchased (Item No. 400000).

Laboratory Procedures

This vial contains lyophilized maturation-inducing steroid (salmonid) (MIS) EIA Antiserum (anti-MIS rabbit IgG). This antiserum has been tested and formulated to work exclusively with ACE™ reagents. This antiserum may not perform as described if used with other assay reagents or protocols. For long term storage, we suggest that the antiserum be stored as supplied at -20°C; it will be stable for at least two years. Reconstitute 100 determination vials with 6 ml EIA Buffer or 500 determination vials with 30 ml EIA Buffer (see buffer preparation instructions below). Store the reconstituted antiserum at 4°C and use within four weeks. For your convenience, we have supplied a 20% surplus of antiserum.

Buffer Preparation

1. Phosphate Buffer

Prepare a 1.0 M phosphate buffer solution by combining 133 g K_2HPO_4 and 32.15 g KH_2PO_4 and diluting to a total volume of 1.0 liter with UltraPure water. The pH of this solution will be 7.4.

2. EIA Buffer

Combine 100 ml of the phosphate buffer prepared above with 100 mg sodium azide, 23.4 g sodium chloride, 370 mg tetrasodium EDTA, and 1 g bovine serum albumin (Sigma A7030 or equivalent). Stir at room temperature until completely dissolved and dilute to a total volume of 1.0 liter with UltraPure water. This buffer may also be purchased as a 10X concentrated buffer (Item No. 400060).

3. Wash Buffer

Combine 10 ml of the 1.0 M phosphate buffer prepared above with 0.5 ml Polysorbate 20. Bring to a final volume of 1.0 liter with UltraPure water. This buffer may also be purchased as a 400X concentrated buffer (Item No. 400062).

Specificity

17 α ,20 β -dihydroxy Progesterone (MIS)	100%	17 α ,20 α -dihydroxy Progesterone	0.04%
4-pregnen-17,20 β ,21-triol-3-one	0.4%	Allopregnanolone	<0.01%
20 β -hydroxy Progesterone	0.1%	Cortisol	<0.01%

Standard Curve

We recommend an eight point standard curve starting at 250 ng/ml. Serially dilute the standard (1:2 dilution) seven times from this point to make the standard curve.

Sensitivity

When used in conjunction with the ACE™ EIA reagents for the MIS (salmonid), this antiserum exhibits a detection limit (80% B/B₀) of ~4 pg/ml and an IC₅₀ (50% B/B₀) of ~15 pg/ml for MIS (salmonid).

Cayman Chemical

Mailing address

1180 E. Ellsworth Road
Ann Arbor, MI
48108 USA

Phone

(800) 364-9897
(734) 971-3335

Fax

(734) 971-3640

E-Mail

custserv@caymanchem.com

Web

www.caymanchem.com

WARNING: THIS PRODUCT IS FOR LABORATORY RESEARCH ONLY: NOT FOR ADMINISTRATION TO HUMANS. NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until information to the contrary becomes available. Do not ingest, swallow, or inhale. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. This information contains some, but not all, of the information required for the safe and proper use of this material. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Cayman Chemical Company makes **no warranty or guarantee** of any kind, whether written or oral, expressed or implied, including without limitation, any warranty of fitness for a particular purpose, suitability and merchantability, which extends beyond the description of the chemicals hereof. Cayman **warrants only** to the original customer that the material will **meet our specifications at the time of delivery**.

Cayman will carry out its delivery obligations with due care and skill. Thus, in no event will Cayman have **any obligation or liability**, whether in tort (including negligence) or in contract, for any direct, indirect, incidental or consequential damages, even if Cayman is informed about their possible existence.

This limitation of liability does not apply in the case of intentional acts or negligence of Cayman, its directors or its employees.

Buyer's **exclusive remedy** and Cayman's sole liability hereunder shall be limited to a **refund** of the purchase price, or at Cayman's option, the **replacement**, at no cost to Buyer, of all material that does not meet our specifications.

Said refund or replacement is conditioned on Buyer giving written notice to Cayman within thirty (30) days after arrival of the material at its destination. Failure of Buyer to give said notice within thirty (30) days shall constitute a waiver by Buyer of all claims hereunder with respect to said material.

For further details, please refer to our Warranty and Limitation of Remedy located on our website and in our catalog.

Copyright Cayman Chemical Company, 09/04/2014

Product Information



Suggested Assay Protocol

This antiserum has been tested and formulated to work exclusively with ACE™ reagents. This antiserum may not perform as described if used with other assay reagents or protocols. *NOTE: This is an abbreviated protocol. If you are not familiar with this assay, please contact us for a complete protocol.*

1. Add 100 µl of EIA Buffer to NSB wells and 50 µl to B₀ wells.
2. Add 50 µl of Standard (Item No. 498504) or sample to the appropriate wells.
3. Add 50 µl Tracer (Item No. 498500) to all wells except Blk and TA.
4. Add 50 µl Antiserum to all wells except Blk, TA, and NSB.
5. Incubate overnight at 4°C.
6. Wash the plate five times with Wash Buffer.
7. Add 200 µl Ellman's Reagent to each well.
8. Add 5 µl Tracer to the TA well.
9. Develop for approximately 90 minutes (B₀ = 0.3-1.0 AU).
10. Read absorbance at a wavelength between 405 and 420 nm.

Blk-Blank; **NSB**-Non-specific Binding; **B₀**-Maximum Binding; **TA**-Total Activity; **S1-S8**-Standards; **1-8**-Samples

	1	2	3	4
A	Blk	S1	S1	1
B	Blk	S2	S2	2
C	NSB	S3	S3	3
D	NSB	S4	S4	4
E	B ₀	S5	S5	5
F	B ₀	S6	S6	6
G	B ₀	S7	S7	7
H	TA	S8	S8	8

Related Products

For a list of related products please visit: www.caymanchem.com/catalog/498502

Cayman Chemical

Mailing address

1180 E. Ellsworth Road
Ann Arbor, MI
48108 USA

Phone

(800) 364-9897
(734) 971-3335

Fax

(734) 971-3640

E-Mail

custserv@caymanchem.com

Web

www.caymanchem.com